

Veröffentlichungen

Veröffentlichungen des Stabes der Biologischen Anstalt Helgoland, die anderenorts erschienen sind

Publications by the staff of the Biologische Anstalt Helgoland that have been published elsewhere

- Andres, H. G., 1981. Die Gammaridae (Crustacea: Amphipoda) der Deutschen Antarktischen Expeditionen 1975/76 und 1977/78. 1. Gammaridae, Melphidippidae und Pagetiniidae. – Mitt. hamb. zool. Mus. Inst. 78, 179–196.
- Andres, H. G., 1981: Lysianassidae aus dem Abyssal des Roten Meeres. – Senckenberg. biol. 61, 429–443.
- Anger, K., Dawirs, R. R., Anger, V. & Costlow, J. D., 1981. Effects of early starvation periods on zoeal development of brachyuran crabs. – Biol. Bull. mar. biol. Lab., Woods Hole 161, 199–212.
- Bartsch, I., 1981. Halacaridae (Acari) aus dem Kanal von Moçambique. – Cah. Biol. mar. 22, 35–63.
- Bartsch, I., 1981. Fünf neue Arten der Gattung *Halacarus* (Acari, Halacaridae) aus dem Atlantik. – Zool. Scr. 10, 203–215.
- Ben-Amotz, A. & Rosenthal, H., 1981. Cryopreservation of marine unicellular algae and early life stages of fish for use in mariculture. – Spec. Publ. Europ. Maricult. Soc. 6, 149–162.
- Bolton, J. J., 1981. Community analysis of vertical zonation patterns on a Newfoundland rocky shore. – Aquat. Bot. 10, 299–316.
- Bulnheim, H.-P. & Scholl, A., 1981. Genetic variation between geographic populations of the amphipods *Gammarus zaddachi* and *G. salinus*. – Mar. Biol. 64, 105–115.
- Drebes, G., 1981. Possible resting spores of *Dissodinium pseudolunula* (Dinophyta) and their relation to other taxa. – Br. phycol. J. 16, 207–215.
- Drebes, H. & Schulz, D., 1981. *Anaulus creticus* sp. nov., a new centric diatom from the Mediterranean Sea. – Bacillaria 4, 161–176.
- Gassmann, G., 1981. Chromatographic separation of diasteriomeric isoprenoids for the identification of fossil oil contamination. – Mar. Pollut. Bull. 12, 78–84.
- Greve, W., Sonntag, N. C. & Kartheus, V., 1981. The Systems impact of inverse trophic relations in carnivorous zooplankton. In: Progress in ecological engineering and management by mathematical modelling. Ed. by D. M. Dubois, Cebedoc, Liège, 715–729.
- Greve, W., 1981. Invertebrate predator control in a coastal marine ecosystem: the significance of *Beroe gracilis* (Ctenophora). Kieler Meeresforsch. (Sonderh.) 5, 211–217.

- Greve, W. & Kartheus, V., 1981. Monitoring ecosystems by mathematical information extraction from indicator species. In: Environmental systems analysis and management. Ed. by S. Rinaldi. CLUP, Milano, 97–99.
- Hamann, I., John, H.-Chr. & Mittelstaedt, E., 1981. Hydrography and its effect on fish larvae in the Mauritanian upwelling area. – Deep-Sea Res. 28A, 561–575.
- Hartmann-Schröder, G., 1981. Die Polychaeten der Fahrten 11, 19, 21 und 26 (1967–1972) von F. S. „Meteor“ in das Gebiet zwischen dem Golf von Biscaya und dem Auftriebsgebiet vor Westafrika. – Meteor-Forsch.-Ergebn. (D) 33, 23–36.
- Hartmann-Schröder, G., 1981. The Ragworm *Nereis diversicolor*. Invertebrates of the Wadden Sea. – Mar. Zool. 4, 113–114.
- Hartmann-Schröder, G., 1981. Zur Kenntnis des Eulitorals der australischen Küsten unter besonderer Berücksichtigung der Polychaeten und Ostracoden. T. 6: Die Polychaeten der tropisch-subtropischen Westküste Australiens (zwischen Exmouth im Norden und Cervantes im Süden). – Mitt. hamb. zool. Mus. Inst. 78, 19–96.
- Hartmann-Schröder, G., 1981. Annelida. In: Lehrbuch der speziellen Zoologie. Hrsg. von H.-E. Gruner. Fischer, Stuttgart, 1, 3, 276–469.
- Hourston, A. S. & Rosenthal, H., 1981. Data summaries for viable hatch from Pacific herring eggs deposited at different intensities on a variety of substrates. – Can. Data Rep. Fish. aquat. Sci. 267, 1–56.
- John, H.-Chr., Thiel, H. & Weikert, H., 1981. Distribution and drift of *Branchiostoma* larvae off Mauritania. – Meeresforsch. 28, 216–227.
- Kessler, H., 1981. Die Seewasserversorgungsanlage der Litoralstation List/Sylt. – Meerestechnik 12, 157–162.
- Kracht, R. & Tesch, F.-W., 1981. Progress report on the eel expedition of R. V. „Anton Dohrn“ and R. V. „Friedrich Heincke“ to the Sargasso Sea. 1979. – Envir. Biol. Fish. 6, 311–375.
- Krüger, F., 1981. Eine neue Funktion zur mathematischen Beschreibung von Wachstumsverläufen. – Biol. Zbl. 100, 195–207.
- Lobban, C. S., Weidner, M. & Lüning, K., 1981. Photoperiod effects enzyme activities in the kelp, *Laminaria hyperborea*. – Z. Pflanzenphysiol. 105, 81–83.
- Lüning, K., 1981. Light. In: The biology of seaweeds. Ed. by C. S. Lobban & M. J. Wynne. Blackwell, Oxford. 326–355. (Botanical Monographs. 17.)
- Lüning, K., 1981. Egg release in gametophytes of *Laminaria saccharina*: induction by darkness and inhibition by blue light and U. V. – Br. phycol. J. 16, 379–393.
- Lüning, K., 1981. Photobiology of seaweeds: ecophysiological aspects. – Proc. int. Seaweed Symp. 10, 35–55.
- Lüning, K., 1981. Photomorphogenesis of reproduction in marine macroalgae. – Ber. dt. bot. Ges. 94, 401–417.
- Martens, P., 1981. On the *Acartia* species of the northern Wadden Sea of Sylt. – Kieler Meeresforsch. (Sonderh.) 5, 153–163.
- Moebus, K., 1981. Enumeration of marine bacteriophages. – Baltic mar. Biol. Publs 3, 45–56.
- Müller, D. G., Gassmann, G., Boland, W., Marner, F. & Jaenicke, L., 1981. *Dictyota dichotoma* (Phaeophyceae): Identification of the sperm attractant. – Science, N. Y. 212, 1040–1041.
- Müller, D. G., Marner, F.-J., Boland, W., Jaenicke, L. & Gassmann, G., 1981. Identifica-

- tion of a volatile gamete secretion in *Spermatochnus paradoxus*. – Naturwissenschaften 67, 478.
- Rieper, M. & Flotow, C., 1981. Feeding experiments with bacteria, ciliates and harpacticoid copepods. – Kieler Meeresforsch. (Sonderh.) 5, 370–375.
- Rosenthal, H. & Oren, O. H. (Eds.), 1981. Research on intensive aquaculture. – Spec. Publ. Europ. Maricult. Soc. 6, 1–278.
- Schnack, S. B. & Elbrächter, M., 1981. On the food of calanoid copepods from the northwest African upwelling region. In: Coastal upwelling. Ed. by F. A. Richards. American Geophysical Union, Washington, D. C., 433–439. (Coastal and Estuarine Sciences, 1.)
- Schulz, K., 1981. *Tharybis minor*, sp. n. (Copepoda: Calanoida: Tharybidae) aus dem nordwestafrikanischen Auftriebsgebiet mit Anmerkungen zur Gattung *Tharybis* Sars. – Mitt. hamb. zool. Mus. Inst. 78, 169–177.
- Siebers, D., Hoppenheit, M. & Harms, U., 1981. Mercury influences uptake of amino acids by marine bacteria. – Mar. Ecol. Prog. Ser. 6, 101–113.
- Sperling, K.-R. & Bahr, B., 1981. Determination of heavy metals in sea water and in marine organisms by flameless atomic absorption spectrophotometry. VIII. Correspondence and some possible sources of error in an intercalibration of cadmium. – Z. analyt. Chem. 306, 71–72.
- Sperling, K.-R., Bahr, B., 1981. Übereinstimmung und einige mögliche Ursachen für Abweichungen bei einer Interkalibration der Cd-Bestimmung. – Ärztl. Lab. 27, 248.
- Tesch, F.-W., 1980. Migratory performance and environmental evidence of orientation. In: Environmental physiology of fishes. Ed. by M. A. Ali. Plenum Press, New York, 589–612.
- Uhlig, G., 1981. Microfaunal food organisms for mariculture. – Spec. Publ. Europ. Maricult. Soc. 6, 93–115.
- Weikert, H. & John, H.-Chr., 1981. Experiences with a multiple opening-closing plankton net (after Bé, modified). – J. Plankton Res. 3, 167–176.
- Westernhagen, H. von, Rosenthal, H., Dethlefsen, V., Ernst, W., Harms, U. & Hansen, P.-D., 1981. Bioaccumulating substances and reproductive success in Baltic flounder *Platichthys flesus*. – Aquat. Toxicol. 1, 85–99.