

Taxonomic biodiversity of geniculate coralline red algae (Corallinales, Rhodophyta) from the Macaronesian region: summary and analysis

Edgar F. Rosas-Alquicira · Rafael Riosmena-Rodríguez ·
Julio Afonso-Carrillo · Ana I. Neto

Received: 30 October 2009 / Revised: 17 June 2010 / Accepted: 23 June 2010 / Published online: 10 July 2010
© Springer-Verlag and AWI 2010

Abstract A catalog and critical review of species and infraspecific taxa of non-fossil geniculate coralline red algae (Corallinales, Rhodophyta) previously reported from the Macaronesian region are presented along with an assessment of species diversity in the region. Published records of geniculate coralline algae are included along with comments relating to type material. Within the catalog, taxa are organized alphabetically by genus and within this by final epithet. From the 31 taxa recorded, 4 are based on type collections from Macaronesian localities. The types of most species and infraspecific taxa reported from the region have yet to be re-examined in a modern context, and most Macaronesian records require verification. The biodiversity of Macaronesian geniculate coralline algae may be lower than current information indicates.

Communicated by Inka Bartsch.

E. F. Rosas-Alquicira (✉) · A. I. Neto
CIIMAR (Centro Interdisciplinar de Investigação
Marinha e Ambiental), Rua dos Bragas 289,
4050-123 Porto, Portugal
e-mail: alquicira@uac.pt

E. F. Rosas-Alquicira · A. I. Neto
Grupo de Biología Marina, Departamento de Biología,
Universidade dos Açores, Rua Mãe de Deus 58, Apart. 1422,
9502 Ponta Delgada, São Miguel, Açores, Portugal

E. F. Rosas-Alquicira · R. Riosmena-Rodríguez
Programa de Investigación en Botánica Marina,
Departamento de Biología Marina, Universidad Autónoma de
Baja California Sur, Km. 5.5, Carretera al Sur. Apdo.
Postal 19-B, C.P. 23080 La Paz, México

J. Afonso-Carrillo
Departamento de Biología Vegetal (Botánica),
Universidad de La Laguna, 38271 La Laguna,
Tenerife, Islas Canarias, España

Keywords Analysis · Baseline · Corallinales · Geniculate · Macaronesia · Taxonomy

Introduction


The Macaronesian region *sensu lato* includes the Azores, Madeira and Salvage Islands (Portugal), the Canaries (Spain) and Cape Verde Islands (Fig. 1). The first studies of (non-fossil) coralline geniculate red algae (Corallinales, Rhodophyta) in this region go back to the beginning of the XIX century with the paper of Lamarck (1815: 233, 237), who described *Corallina millegrana* and *Corallina purpurea* for the Canary Islands.

An important contribution to the knowledge of Macaronesian Corallinales, with an extensive collection for the region, was done by the Dutch oceanographic expeditions CANCAP, in the 1970 and 1980s, with material collected in the Azores, Madeira, Salvage, the Canaries and Cape Verde Islands (Prud'homme van Reine 1988).

Historical revisions of phycological studies including the coralline algae are those by Neto (1994) for the Azores, Levring (1974) and Neto et al. (2001) for Madeira, Afonso-Carrillo (1988) and Haroun et al. (2002) for the Canary Islands and Prud'homme van Reine (1988) for Cape Verde Islands. The accounts of the coralline algae published by Price et al. (1986, 1992) include information on Macaronesian material but only for Cape Verde and the Canary Islands. The work of John et al. (2004) also included data from Madeira but the Azores were not considered in any of these publications.

Currently, there are 31 names of geniculate coralline seaweeds reported for the Macaronesian region but there is no published information compiling the data from the various archipelagos. Information is also lacking on basionym,

Fig. 1 The Macaronesian region



type material locality, localization, and illustrations, current nomenclatural situation and taxonomic status.

The present study provides a baseline summary and analysis of existing taxonomic information on the geniculate Corallinales for the Macaronesian region *sensu lato*, along with comments and/or nomenclatural notes as required.

Arrangement of the catalog

The species records were compiled from identification keys, floristic studies, checklists and published catalogs. Unpublished theses and reports were not considered because some involved names that are not effectively published in the context of the International Code of Botanical Nomenclature (ICBN, McNeill et al. 2006).

The type material of a number of taxa of geniculate Corallinales reported from the Macaronesian region has not been studied in the modern context of taxonomy, meaning that the descriptions are incomplete and factually inaccurate, include criteria that are unreliable and are not based on original research encompassing specific morphological, anatomical and ultrastructural investigations. The presence or absence of genicula in those taxa has been inferred from the genus in which the taxon was placed originally. Thus, taxa whose types have not been studied in a modern context but that originally were described as species or infraspecific taxa of *Amphiroa* J. V. Lamouroux, *Cheilosporum* (Decaisne) Zanardini, *Corallina* Linnaeus, *Haliptilon* (Decaisne)

Lindley and *Jania* J. V. Lamouroux are presumed to possess genicula.

Taxa are listed in alphabetical order by the currently accepted name for the genera and constituent species. For each taxon, the following information is provided: basionym, type material localization, type locality, published illustrations of type material, reference to its study in a modern context and current placement/name. Table 1 summarizes this information. Published records for the Macaronesian region are also provided, following a north–south latitudinal order: Azores, Madeira, Salvage Islands, Canary Islands and Cape Verde Islands. For each archipelago, homotypic synonyms are arranged in chronological order, with author citations and publications details provided for each.

In the case of taxa for which the type material is not known, or when known, was not revised, nor were the specimens studied in the modern context of taxonomy the status and/or disposition of the taxa is reported as uncertain. Family names of authors are written out in full and Herbarium abbreviations follow Holmgren and Holmgren (1998).

Catalog

Amphiroa

Currently, the genus *Amphiroa* is recognized as one of six genera in the Corallinaceae, subfamily Lithophylloideae (Bailey 1999, Table 4, p. 214). Presently its generic concept

Table 1 Current status and disposition of geniculate Corallinales reported from the Macaronesian region

Current placement/name	Basionym	Valid publication	Type material studied in modern context	Remarks	Species status
Genus	Epithet				
<i>Amphiroa</i>	<i>beauvoisi</i>	= Lamouroux 1816: 299–300	Lamouroux 1816: 299–300	Norris and Johansen 1981, Harvey et al. 2009	Distinct species
<i>cryptarthrodia</i>	<i>Coralina verrucosa</i>	Zanardini 1840: 136	Harvey 1849: 95	Rosas-Alquicira et al. 2010	Distinct species
<i>exilis</i>	= <i>Coralina fragilissima</i>	Linnaeus 1758: 806	Weber-van Bosse 1904	Uncertain	Uncertain
<i>fragilissima</i>	= <i>rigida</i>	Lamouroux 1816: 297	Norris and Johansen 1981	Hamel and Lemoine 1953,	Distinct species
<i>rigida</i>	= <i>Amphiroa verrucosa</i>	Kützing 1841: 18	Hamel and Lemoine 1953, Rosas-Alquicira et al. 2010	Synonym of A. <i>cryptarthrodia</i>	Synonym of
<i>Chiilosporum</i>	<i>elegans</i>	<i>Amphiroa elegans</i>	Generic disposition as <i>Cheiilosporum</i> certain	Uncertain	Uncertain
<i>Corallina</i>	<i>deshayesii</i>	=	Designation/localization of the type material is required	Uncertain	Uncertain
<i>elongata</i>	=	Hooker and Harvey 1849; 101	Irvine and Johansen 1994	Designation/localization of the type material is required	Distinct species
<i>granifera</i>	=	Montagne 1846: 130	Only a fragment of the holotype was studied (Garbarey and Johansen 1982)	Uncertain	Uncertain
<i>lobata</i> ^a	=	Lamouroux 1816: 286	Generic disposition as <i>Corallina</i> certain	Uncertain	Uncertain
<i>mediterranea</i>	=	Araschouq in Agardh 1852: 568	Considered a synonym of <i>C. elongata</i> (Irvine and Johansen 1994), but without comparing both types	Uncertain	Uncertain
<i>microptera</i> ^a	=	Montagne 1846: 130	Only a fragment of the holotype was studied (Garbarey and Johansen 1982)	Generic disposition as <i>Corallina</i> certain	Uncertain
<i>millegrana</i> ^a	=	Lamarck 1815: 233	Only a fragment of the holotype was studied (Garbarey and Johansen 1982)	Generic disposition as <i>Corallina</i> certain	Uncertain
<i>officinalis</i>	=	Linnaeus 1758: 805	Irvine and Johansen 1944, Walker et al. 2009	Distinct species	Distinct species
<i>Halipitilon</i>	<i>cubensis</i>	Jania cubensis	Montagne in Kützing 1849: 709–710	Generic disposition as <i>Halipitilon</i> certain	Uncertain
<i>purpuratum</i> ^a	<i>Coralina purpurata</i>	Lamarck 1815: 237	Only a fragment of the holotype was studied (Garbarey and Johansen 1982)	Generic disposition as <i>Halipitilon</i> certain	Uncertain

Table 1 continued

Current placement/name		Basionym	Valid publication	Type material studied in modern context	Remarks	Species status
Genus	Epithet					
<i>squamatum</i>	<i>Corallina squamata</i>	Linnaeus 1758: 806	Johansen et al. 1973, Irvine and Johansen 1994	No	Distinct species	
<i>virgatum</i>	<i>Corallina virgata</i>	Zanardini 1844: 1025	Only a fragment of the type was studied (Garibay and Johansen 1982)	Generic disposition as <i>Haliptilon</i> certain	Uncertain	
<i>Jania adhaerens</i>	=	Lamouroux 1816: 270			Uncertain	
<i>capillacea</i>	=	Harvey 1853: 84–85			Uncertain	
<i>crassa</i>	=	Lamouroux 1821: 23, Johansen and Womersley 1944			Uncertain	
<i>intermedia</i>	<i>Corallina intermedia</i>	Kützing 1858: 37–38, 42		Designation/localization of the type material is required	<i>Nomen nudum</i> (Woelkerling and Nelson 2004)	
<i>longifurca</i>	=	Zanardini 1844: 1025		Designation/localization of the type material is required	Uncertain	
<i>micrarthrodia</i>	=	Lamouroux 1816: 271	Johansen and Womersley 1944	No	Distinct species	
<i>natalensis</i>	=	Harvey 1849: 107		Considered a synonym of <i>Jania verrucosa</i> (Johansen and Womersley 1944), but without comparing both types	Uncertain	
<i>pumila</i>	=	Lamouroux 1816: 269		Holotype in poor condition being difficult to interpret (Johansen and Womersley 1944)	Uncertain	
<i>nubens</i> var. <i>corniculata</i>	<i>Corallina corniculata</i>	Linnaeus 1758: 806			Uncertain	
<i>nubens</i> var. <i>rubens</i>	<i>Corallina rubens</i>	Linnaeus 1758: 806	Irvine and Johansen 1944	Distinct species		
<i>tenella</i>	<i>Corallina tenella</i>	Kützing 1858: 41			Uncertain	
<i>verrucosa</i>	=	Lamouroux 1816: 270		Considered a distinct species of <i>Jania</i> (Johansen and Womersley 1944) but without studying its type	Uncertain	

^a Species based on types from the Macaronesian region

is consensual and a detailed diagnosis is provided by Womersley and Johansen (1996: 283–284) and Harvey et al. (2009: 259; Table 1, p. 260). The relationship between *Amphiroa* and other Lithophylloideae genera is provided in Woelkerling et al. (2002, Table 2, p. 370; Table 3, p. 372). The work by Afonso-Carrillo and Sansón (1999: 33–34) provides a key for species diagnosis in the Canaries.

Amphiroa beauvoisii J. V. Lamouroux

Basionym: *Amphiroa beauvoisii* J. V. Lamouroux 1816: 299–300. **Holotype:** CN (Norris and Johansen 1981: 6); illustrations: Norris and Johansen (1981, Fig. 7b, p. 6) and Harvey et al. (2009, Figs. 18–21, p. 270; Figs. 22–23, p. 271). **Type locality:** Portugal coasts (Lamouroux 1816: 299).

Current placement/name: *A. beauvoisii* J. V. Lamouroux according to Norris and Johansen (1981: 6) and Harvey et al. (2009: 268–273) based on the study of the holotype.

Published records:

AZORES: Gain (1914: 22), Schmidt (1931: 67), Feldmann (1946: 421), Ardré (1970: 224), Cordeiro-Marino (1978: 46), South and Tittley (1986: 42), Reyes and Sansón (1991: 74), Schneider and Searles (1991: 226), Neto (1994: 25), Prud'homme van Reine et al. (1994: 70), Tittley and Neto (1994: 7), Cremades et al. (1997: 13), and Rosas-Alquicira and Neto (personal observation).

MADEIRA: Levring (1974: 68), Augier (1985: 101) and John et al. (2004: 58).

SALVAGE ISLANDS: John et al. (2004: 58).

CANARY ISLANDS: Afonso-Carrillo et al. (1984: 30), Price et al. (1986: 10), Gil-Rodríguez et al. (1987: 146, 2003: 25), Viera-Rodríguez (1987: 241), Afonso-Carrillo and Sansón (1989: 26; 1999: 195), Guadalupe et al. (1995: 37), Haroun et al. (2002: 146), Sangil et al. (2003a: 87; 2004: 89) and Muñoz et al. (2007: 107).

Amphiroa cryptarthrodia Zanardini

Basionym: *Corallina verrucosa* Zanardini 1840: 136. **Neotype:** MCVE unnumbered, Zanardini's original collection nr. 14 (Rosas-Alquicira et al. 2010); illustrations: Hamel and Lemoine (1953, pl. 5, p. 43) and Rosas-Alquicira et al. (2010, Figs. 1, 4–20) as *Amphiroa verruculosa* Kützing; Rosas-Alquicira et al. (2010, Figs. 27–36) as *Amphiroa cryptarthrodia* Zanardini. **Syntype localities:** Trieste and Dalmazia, Adriatic Sea (Rosas-Alquicira et al. 2010).

Current placement/name: *A. cryptarthrodia* Zanardini according to Rosas-Alquicira et al. (2010) based on the

study of neotype material. Afonso-Carrillo et al. (1983: 46) cited the presence of *A. cryptarthrodia* at the Canaries as dubious.

Published records:

AZORES: Schmidt (1931: 67), Audiffred and Weisscher (1984: 17), South and Tittley (1986: 42), Neto (1994: 25), Tittley and Neto (1994: 7) and Rosas-Alquicira and Neto (personal observation).

MADEIRA: Levring (1974: 68), Audiffred and Weisscher (1984: 17, 33) and John et al. (2004: 58).

SALVAGE ISLANDS: Audiffred and Weisscher (1984: 17, 33), Price et al. (1986: 11) and John et al. (2004: 58).

CANARY ISLANDS: Lemoine (1929: 71), Gil-Rodríguez and Afonso-Carrillo (1980: 35), Price et al. (1986: 11) and John et al. (2004: 58).

Amphiroa exilis Harvey

Basionym: *Amphiroa exilis* Harvey 1849: 95. **Syntypes:** TCD specimens Darwin 595 (with annotation Det. H.W. Johansen Sept 1967) and Darwin 629 (two specimens), BM Darwin 595 (one sheet with the stamp of Hookerianum Herbarium 1867 and the annotation *A. exilis*) and Darwin 629 (with the annotation *A. exilis*); illustrations: Porter (1987, BM Darwin 595, Fig. 1, p. 188; BM Darwin 629, Fig. 2, p. 189). **Type locality:** Botofogo Bay, Rio de Janeiro, Brazil (Porter 1987: 188–189); Algoa Bay, Cape Province, South Africa (Silva et al. 1996: 220).

Current placement/name: *A. beauvoisii* J. V. Lamouroux but status and disposition uncertain as the type was not studied in a modern context. Yendo (1905: 4) based on Madam Weber-van Bosse observations (1904: 100) of Harvey's plants (*A. exilis* presumed type material) proposed *A. exilis* as a heterotypic synonym of *A. beauvoisii*. The synonym *A. beauvoisii* was accepted by Prud'Homme van Reine (1994: 70) based on the concept of Price et al. (1986) for the Azorean material of *A. exilis* that had been collected and identified by Piccone (1889). In contrast to Prud'Homme van Reine et al. (1994), we think the name *A. exilis* should be kept until more comparisons and study of type material have been conducted.

Published records:

AZORES: Agardh (1870: 363), Piccone (1889: 209) and Tittley et al. (1998: 467).

Amphiroa fragilissima (Linnaeus) J. V. Lamouroux

Basionym: *Corallina fragilissima* Linnaeus 1758: 806. **Holotype:** LINN (Yoshida 1998: 533); illustrations:

Lamouroux (1821, Table. 21, Fig. d) and Weber-van Bosse (1904, Fig. 19, pl. XVI) as *Amphiroa fragilissima* (Linnaeus) J. V. Lamouroux. *Type locality*: Jamaica (Silva et al. 1996: 223).

Current placement/name: *A. fragilissima* (Linnaeus) J. V. Lamouroux according to Weber-van Bosse (1904: 90–91) based on the study of type material.

Published records:

AZORES: *Amphiroa fragilissima* (Linnaeus) J. V. Lamouroux. South and Tittley (1986: 42), Prud'homme van Reine (1988: 179), Neto (1994: 25) and Rosas-Alquicira and Neto (personal observation).

MADEIRA: *Amphiroa fragilissima* (Linnaeus) J. V. Lamouroux. Levring (1974: 69).

Amphiroa? fragilissima (Linné) J. V. Lamouroux. Neto et al. (2001: 403).

Amphiroa fragilissima. Grunow (1868: 79).

CANARY ISLANDS: *Amphiroa fragilissima* J. V. Lamouroux. Afonso-Carrillo et al. (1983: 30), Prud'homme van Reine et al. (1994: 70), Haroun et al. (2002: 146).

Amphiroa fragilissima (Linnaeus) J. V. Lamouroux. Afonso-Carrillo (1980b: 54), Afonso-Carrillo et al. (1984: 32), Price et al. (1986: 73), Afonso-Carrillo (1986: 190), González et al. (1986: 315), Viera-Rodríguez (1987: 242), Reyes and Sansón (1991: 74), Elejabeitia et al. (1992: 3), Pinedo et al. (1992: 32), Guadalupe et al. (1995: 37), Afonso-Carrillo and Sansón (1999: 195), Gil-Rodríguez et al. (2003: 25), Montañés et al. (2003: 124) and Sangil et al. (2003a: 87; b: 310; 2004: 89).

Amphiroa fragilissima. Jorge et al. (1984: 118) and Afonso-Carrillo and Sansón (1989: 26).

CAPE VERDE ISLANDS: *Amphiroa fragilissima* J. V. Lamouroux. Price et al. (1986: 73).

Amphiroa fragilissima (Linnaeus) J. V. Lamouroux. Price et al. (1986: 73) and Otero-Schmitt and Sanjuan (1992: 382).

Amphiora rigida J. V. Lamouroux

Basionym: *Amphiroa rigida* J. V. Lamouroux 1816: 297.

Holotype: CN (Norris and Johansen 1981: 20); illustrations: Lamouroux (1816, pl. 11, Fig. 1, p. 297); Cremades et al. (1997, PC, Fig. C, p. 16). *Type locality*: Mediterranean (Lamouroux 1816: 297).

Current placement/name: *A. rigida* J. V. Lamouroux according to Norris and Johansen (1981: 21) based on study of holotype material. The presence of this species in the

Canary Islands was confirmed by Haroun et al. (2002: 139) but authors did not mention if they saw the material. This species is closely related to *A. cryptarthrodia* Zanardini, and it is likely that the reports of *A. rigida* in the Azores correspond to misdeterminations of *A. cryptarthrodia* (Rosas-Alquicira and Neto, personal observation).

Published records:

AZORES: Piccone (1889: 208), South and Tittley (1986: 42), Neto (1994: 25) and Tittley et al. (1998: 467).

Amphiroa cf. rigida J.V. Lamouroux. Tittley and Neto (1994: 7).

MADEIRA: Levring (1974: 68) and John et al. (2004: 59).

CANARY ISLANDS: Price et al. (1986: 12), Guadalupe et al. (1995: 37), Gil-Rodríguez et al. (2003: 25) and John et al. (2004: 59).

Amphiroa rigida. Afonso-Carrillo and Sansón (1989: 26).

CAPE VERDE ISLANDS: John et al. (2004: 59).

Amphiroa verruculosa Kützing

Basionym: *Amphiroa verrucosa* Kützing 1841: 18. *Holotype*: L 0055675; illustrations: Kützing (1849, pl. 79, fig. III, p. 700) and Rosas-Alquicira et al. (2010, Figs. 27–36) as *Amphiroa verruculosa*. *Type locality*: Adriatic Sea (Kützing 1849: 700).

Current placement/name: *A. cryptarthrodia* Zanardini according to Rosas-Alquicira et al. (2010) based on the study of neotype material. Neto et al. (2001: 412) cited a doubtful record of *A. verruculosa* from the Archipelago of Madeira.

Published records:

MADEIRA: Feldmann (1946: 425) and Neto et al. (2001: 412).

Cheilosporum

The genera *Cheilosporum*, *Haliptilon* and *Jania* were grouped into the Janieae tribe, subfamily Corallinoideae, based on the combination of the following features: (1) presence of a thick carposporophytic fusion cell; (2) gonimoblastic filaments arising from fusion cell margins; (3) male conceptacles chambers narrow with a short canal and (4) few tetrasporangia per conceptacle (Johansen and Silva 1978: 413; Table 1, p. 414). According to these authors, *Cheilosporum* distinguishes from other Janieae genera by possessing a marginal rather than axial location of conceptacles and the presence of pronounced intergenicular lobes.

A complete diagnosis of the genus was provided by Johansen (1981, Appendix 1, p. 216). Womersley and Johansen (1996: 296) included a key for the Janiae genera in their book for Southern Australia where they mentioned as segregating generic characters, the branching pattern of the main axes, the intergeniculate shape and the conceptacles position. Kim et al. (2007), based on combined molecular (SSU rDNA) and anatomical data (Table 1, p. 1313), determined that these characters were not useful to separate the three genera. They concluded that the Janiae should be merged only into the single genus *Jania* recognized by the generic characters previously listed by Johansen and Silva (1978: 413; Table 1, p. 414). This conclusion was followed by Woelkerling et al. (2008). Walker et al. (2009), based on cox1 and 18S rRNA analysis of *Haliptilon squamatum* and *Jania rubens*, also concluded they were not distinct but suggested that the clarification of the phylogenetic position of *Haliptilon* and *Jania* required more comparisons and study of original material of the genera involved. We share the opinion of Walker et al. (2009), a reason why in the present catalog the genera and species of *Cheilosporum*, *Haliptilon* and *Jania* are treated separately.

Cheilosporum elegans J. E. Areschoug

Basionym: *Amphiroa elegans* J. D. Hooker and Harvey 1849: 101. **Type:** TCD Colenso specimen 630; illustrations: Harvey (1849, pl. 38 lower left, p. 101). **Type locality:** New Zealand (Woelkerling and Nelson 2004: 54).

Current placement/name: *Cheilosporum elegans* J. E. Areschoug but status and disposition of species uncertain as the type was not studied in a modern context (Woelkerling and Nelson 2004: 55). This conclusion was based on specimens originally described by Hooker and Harvey (in Harvey 1849: 101) under the name *Amphiroa elegans*, an illegitimate name (ICBN Art. 53.1) since it is a later homonym of *Amphiroa elegans* Sonder (1845: 55), now considered a heterotypic synonym of *Metagoniolithon stelliferum* (Lamarck) Ducker (Silva et al. 1996: 260). Womersley and Johansen (1996: 315, 317) mentioned *C. elegans* J. E. Areschoug and also *Amphiroa elegans* J. D. Hooker & Harvey in Harvey, *nom. illeg.* as heterotypic synonyms of *Cheilosporum sagittatum* (Lamouroux) J. E. Areschoug. A similar suggestion was made by Moura and Guimarães (2002: 69) for *C. elegans* (J. D. Hooker & Harvey in Harvey) Areschoug. These suggestions, however, were made without reviewing type material. Later, Kim et al. (2007) based on molecular and reproductive features proposed to merge material of *Cheilosporum* and *Haliptilon* within *Jania* genus and suggested the new combination *Jania cultrata* (Harvey) J. H. Kim, Guiry & H.-G. Choi comb. nov. This conclusion, however, was not based on type material observations.

Published records:

CAPE VERDE ISLANDS: *Cheilosporum elegans* (J. D. Hooker & Harvey) J. E. Areschoug. Askenasy (1896: 26) and Price et al. (1986: 65). *Cheilosporum elegans* Harvey. Price et al. (1986: 65).

Corallina

Currently, the genus *Corallina* is recognized to belong to the Corallinaceae, subfamily Corallinoideae (Johansen 1981, Table 9, p. 188; Harvey et al. 2003, Table 4, p. 995). Its generic concept is consensual. The main characteristics are listed in Johansen (1981: 217), and a detailed diagnosis is provided by Womersley and Johansen (1996: 289–290). The relationship between *Corallina* and other Corallinoideae genera is provided by Woelkerling et al. (2008, Table 4, p. 282; Table 5, p. 287).

Corallina deshayesii Montagne

Basionym: *Corallina deshayesii* Montagne 1846: 130. **Type:** not found; illustrations: not found. **Type locality:** not found.

Current placement/name: *C. deshayesii* Montagne but status and disposition uncertain as the type was not studied in a modern context. Price et al. (1986: 73) mentioned that *C. deshayesii* may well be a synonym of *Corallina mediterranea* J. E. Areschoug. The designation/localization of the type material and its study in a modern context is required.

Published records:

SALVAGE ISLANDS: Price et al. (1986: 73).

Corallina elongata J. Ellis & Solander

Basionym: *Corallina elongata* J. Ellis and Solander 1786: 119. **Lectotype:** Ellis's illustrations designated by Irvine and Johansen (1994: 41); illustrations: Ellis (1755, pl. 24, Fig. 3, p. 49). **Type locality:** Cornwall, England (Irvine and Johansen 1994: 41).

Current placement/name: *C. elongata* J. Ellis & Solander according to Irvine and Johansen (1994: 41) based on the study of lectotype material. Walker et al. (2009) in their integrative taxonomic approach including phylogenetic reconstructions based on the cox1 and 18S rRNA genes in addition to a comparative morphological assessment study identified three clusters. *C. officinalis* *sensu stricto* and *C. elongata* from Britain corresponded to similar groups in the cox1 tree whereas the sequences of the unidentified *Corallina* specimens from Madeira and Tenerife clustered with sequences of *C. caespitosa* R. H. Walker, J. Brodie &

L. M. Irvine and *C. elongata* from other locations. The results indicate that the name *C. elongata* has been misapplied in both the Atlantic and Mediterranean. While the concept of the species is not disputed for those samples from Britain and Ireland, the samples from Greece are *C. caespitosa*, and those from elsewhere in the Atlantic are of an unknown species and should be re-evaluated. Higher divergences as observed in the sequences identified to the genus from Tenerife and Madeira in the cox1 gene can be interpreted as the result of geographical isolation and gene pool fragmentation (Hebert et al. 2003). These islands are geographically isolated in the Atlantic but it is not known if the populations interbreed with populations from elsewhere in the Atlantic and Mediterranean. For the relationship between *C. elongata* and *C. mediterranea* J. E. Areschoug, see *C. mediterranea* entry.

Published records:

AZORES: Castro and Viegas (1983: 20), South and Tittley (1986: 42), Tittley and Neto (1994: 8), Toste et al. (2003: 1270) and Wallenstein and Neto (2006: 199).

Corallina elongata Johnst. Castro and Viegas (1987: 63).

Corallina elongata. Neto (1994: 26, 2000a: 139, b: 490) and Prud'homme van Reine et al. (1994: 81) and Tittley and Neto (2000: 20).

MADEIRA: Audiffred and Prud'homme van Reine (1985: 38) and John et al. (2004: 73).

Corallina elongata. Prud'homme van Reine et al. (1994: 81).

SALVAGE ISLANDS: Price et al. (1986: 73), Prud'homme van Reine et al. (1994: 81) and John et al. (2004: 73).

CANARY ISLANDS: Price et al. (1986: 73), Viera-Rodríguez and Wildpret de la Torre (1986: 215), González et al. (1986: 316), Viera-Rodríguez (1987: 242), Viera-Rodríguez et al. (1987: 273), Reyes and Sansón (1991: 79), Ballesteros et al. (1992: 516), Pinedo et al. (1992: 36), Kristiansen et al. (1993: 95), Irvine and Johansen (1994: 42), Prud'homme van Reine et al. (1994: 81), Guadalupe et al. (1995: 37), Rojas-González and Afonso-Carrillo (2000: 129, 2004: 138), Gil-Rodríguez et al. (2003: 25), Haroun et al. (2003: 108), Montañés et al. (2003: 124), Sangil et al. (2003a: 94 103, b: 311, 2004: 89, 2005: 326), Afonso-Carrillo and Sobrino (2004: 149), John et al. (2004: 73), Muñoz et al. (2007: 107) and Afonso-Carrillo et al. (2009: 238).

Corallina elongata. Afonso-Carrillo (1980a: 7, b: 54), Delgado et al. (1984: 105), Haroun et al. (1984: 111), Jorge et al. (1984: 118), Afonso-Carrillo and Sansón (1989: 25), Gil-Rodríguez et al. (1992: 101), Hardisson et al. (1998: 945-950) and Tuya and Haroun (2006: 17).

Corallina granifera J. Ellis & Solander

Basionym: *Corallina granifera* J. Ellis & Solander 1786: 120. **Type:** whereabouts uncertain (Woelkerling and Nelson 2004: 56); illustrations: Ellis and Solander (1786, Table 21, Fig. c.C., p. 120). **Type locality:** Mediterranean coasts of Africa (Lamouroux 1816: 288).

Current placement/name: *C. granifera* J. Ellis & Solander but status and disposition uncertain as the type was not studied in a modern context. Ardisson (1883: 465) and Yendo (1905: 30) cited *C. granifera* as a synonym of *Corallina virgata* Zanardini but no revision of type material was made. The designation/localization of the type material and its study in a modern context is required.

Published records:

AZORES: Gain (1914: 22), Lemoine (1924: 130), Schmidt (1929b: 172, 1931: 65), Feldmann (1946: 421), Weisscher (1982: 31, 1983: 63) and Audiffred and Weisscher (1984: 20).

Corallina granifera. Chapman (1955: 801) and Audiffred and Prud'homme van Reine (1985: 38).

CANARY ISLANDS: Price et al. (1986: 74) and González et al. (1986: 316).

Corallina granifera. Afonso-Carrillo (1980b: 54), Delgado et al. (1984: 105) and Price et al. (1986: 74).

Corallina granifera J. Ellis & Solander?. Ardré (1970: 228).

Jania granifera Decaisne. Vickers (1896: 306) and Price et al. (1986: 74).

MADEIRA: Lemoine (1924: 130), Børgesen (1929: 70), Schmidt (1931: 101), Seoane-Camba (1965: 110), Weisscher (1982: 31, 1983: 63), Audiffred and Weisscher (1984: 20, 33) and Audiffred and Prud'homme van Reine (1985: 38).

Corallina granifera J. Ellis & Solander?. Ardré (1970: 228).

Jania granifera Decaisne. Gain (1914: 7) and Neto et al. (2001: 403).

SALVAGE ISLANDS: Price et al. (1986: 74).

CAPE VERDE ISLANDS: Weisscher (1982: 31) and Price et al. (1986: 74).

Corallina lobata J. V. Lamouroux

Basionym: *Corallina lobata* J. V. Lamouroux 1816: 286.

Holotype: CN; illustrations: not found. **Type locality:** The Canaries (Lamouroux 1816: 286).

Current placement/name: *C. lobata* J. V. Lamouroux. Examination of a fragment of the type material revealed stable characters at the genus level (Garbarek and Johansen 1982: 212), but the status as a distinct species within the

genus remains uncertain until comparative studies of relevant types are undertaken. Afonso-Carrillo (unpublished information) verified that the type material of *C. lobata* was not different from the actual concept of *Haliptilon virgatum* (Zanardini) Garbary & H. W. Johansen (for further comments see Price et al. 1992: 125). As *C. lobata* is an older name, it would have priority over *C. virgata* Zanardini (1840) (Afonso-Carrillo and Sansón 1999: 151–152). For more information, see *Haliptilon virgatum* entry.

Published records:

- MADEIRA: Afonso-Carrillo and Sansón (1999: 151). SALVAGE ISLANDS: Afonso-Carrillo and Sansón (1999: 151). CANARY ISLANDS: Lamouroux (1816: 286), Afonso-Carrillo et al. (1983: 46, 1984: 26, 30), Price et al. (1986: 75), Reyes and Afonso-Carrillo (1993: 127), Afonso-Carrillo and Sansón (1999: 151). *Corallina lobata*. Delgado et al. (1984: 105). CAPE VERDE ISLANDS: Afonso-Carrillo and Sansón (1999: 151).

Corallina mediterranea J. E. Areschoug

Basionym: *Corallina mediterranea* J. E. Areschoug 1852: 568. *Type*: Herb. Reg. Acad. Scientiarum Holm; illustrations: not found. *Type locality*: Mediterranean (Agardh 1852: 568).

Current placement/name: *C. mediterranea* J. E. Areschoug but status and disposition uncertain as the type was not studied in a modern context. Ardré (1970: 226) and Solms-Laubach (1881 in Ardisson 1883: 464) mentioned that the criteria used to distinguish *C. mediterranea* and *Corallina officinalis* Linnaeus were unsatisfactory and that further research was needed before any decision to merge both species. The historical use of *C. mediterranea* or *C. officinalis* at Salvagens Islands (Madeira) and in the Canary Islands is discussed in Price et al. (1986: 74).

Prud'homme van Reine et al. (1994: 81) redefined the *C. mediterranea* material recorded by Grunow (1868: 77) for the Archipelago of Madeira as *C. elongata* J. Ellis & Solander in the Afonso-Carrillo and Sansón (1989) concept. This synonym was generalized by Price et al. (1986: 73, as *C. elongata* Johnst. Newton 1931: 313) and published as valid by Irvine and Johansen (1994: 41) but only based on the study of the lectotype of *C. elongata*.

For details on the relationship between *C. mediterranea*, *C. elongata* and *C. officinalis*, see *C. mediterranea* and *C. officinalis* entries.

Published records:

- AZORES: Lemoine (1924: 130), Schmidt (1931: 65), Feldmann (1946: 419), Ardré et al. (1974: 178), Tittley et al. (1998: 468) and Neto (2000b: 485, 2001: 104). MADEIRA: Grunow (1868: 77), Lemoine (1924: 130), Schmidt (1931: 101) and John et al. (2004: 73). SALVAGE ISLANDS: Price et al. (1986: 73). CANARY ISLANDS: Lemoine (1924: 130), Børgesen (1929: 68), Schmidt (1931: 101), Feldmann (1946: 407), Seoane-Camba (1965: 110), Kohlmeyer (1967: 143), Santos Guerra et al. (1970: 24), Acuña González (1972: 5), Santos Guerra (1972: 95), Perez-Cirera (1975: 19), Afonso-Carrillo et al. (1978: 232) and Price et al. (1986: 73). *Corallina mediterranea*. Vickers (1896: 295), Sauvageau (1912: 210) and González (1977b: 100). *Corallina mediterranea* L. Gil-Rodríguez (1980: 138).

Corallina microptera Montagne

Basionym: *Corallina microptera* Montagne 1846: 130. *Type*: M Bourgeau; PC (Afonso-Carrillo and Sansón 1999: 152); illustrations: not found. *Type locality*: Orotava, Tenerife, the Canaries (Montagne 1846: 130).

Current placement/name: *C. microptera* Montagne. Examination of a fragment of the type material revealed stable characters at the genus level (Garbary and Johansen 1982: 212), but the status as a distinct species within genus remains uncertain until comparative studies of relevant types are undertaken. According to Bornet (1892), *C. microptera* from the Canary Islands is just a small form of *C. elongata* (as *C. mediterranea* J. E. Areschoug, Afonso-Carrillo and Sansón 1999: 152). Revision of the type material revealed stable characters at the genus level (Garbary and Johansen 1982: 212). The study of type specimens in a modern context is required for specific diagnosis.

Published records:

- AZORES: *Corallina microptera*. Afonso-Carrillo and Sansón (1999: 152). MADEIRA: Afonso-Carrillo and Sansón (1999: 152). CANARY ISLANDS: Montagne (1846: 130).

Corallina millegrana Lamarck

Basionym: *Corallina millegrana* Lamarck 1815: 233. *Holotype*: PC 0028655 General Herbarium; illustrations: not found. *Type locality*: Tenerife, the Canaries (Lamarck 1815: 233).

Current placement/name: *C. millegrana* Lamarck. Examination of a fragment of the type material revealed stable

characters at the genus level (Garbary and Johansen 1982: 212), but the status as a distinct species within genus remains uncertain until comparative studies of relevant types are done.

Published records:

CANARY ISLANDS: Afonso-Carrillo and Sansón (1999: 152).

Corallina officinalis Linnaeus

Basionym: *Corallina officinalis* Linnaeus 1758: 805. *Lectotype:* LINN 1293. 9 (Irvine and Johansen 1994: 44); illustrations: Walker et al. (Walker et al. 2009, C. officinalis, Fig. 5a; *Corallina compacta* Crouan & Crouan (PC), Fig. 5b; *Corallina nana* Zanardini (MCVE), Fig. 5c; *Corallina calvadosii* J. V. Lamouroux (PC), Figs. 5 d–e, p. 296). *Type locality:* European Seas (Lamouroux 1816: 284).

Current placement/name: *C. officinalis* Linnaeus according to Irvine and Johansen (1994: 44) and Walker et al. (2009: 295) based on the study of lectotype material. Afonso-Carrillo et al. (1983: 46) mentioned the presence of *C. officinalis* in the Canary Islands as doubtful. Prud'homme van Reine et al. (1994: 81) studying the material collected by Piccone and housed at PAD Herbarium redefined *C. officinalis* from Azores (Piccone 1889: 209), Canaries (Piccone 1884: 42) and Madeira (Piccone 1884: 42) as *C. elongata* J. Ellis & Solander in the concept of Afonso-Carrillo (personal communication). However, no revision of the type material was made. Irvine and Johansen (1994: 44) provided a detailed account of *C. officinalis* in the British Isles while Womersley and Johansen (1996: 291) provided an account of the species in southern Australia.

For details on the relationship between *C. officinalis*, *C. elongata* and *C. mediterranea* J. E. Areschoug, see *C. elongata* and *C. mediterranea* entries.

Published records:

AZORES: Piccone (1889: 209), Gain (1914: 15, 22), Schmidt (1931: 65), Feldmann (1946: 419), Levring (1974: 69), Audiffred (1985: 174), Weisscher (1983: 64), Audiffred and Prud'homme van Reine (1985: 38), South and Tittley (1986: 42), Neto (1994: 26), Tittley and Neto (1994: 8, 2000: 20) and Tittley et al. (1998: 468).

Corallina officinalis. Schmidt (1929a: 331).

MADEIRA: Piccone (1884: 42–43), Barton (1897: 374), Gain (1914: 7), Schmidt (1931: 101), Levring (1974: 69), Weisscher (1983: 64), Audiffred and Weisscher (1984: 33), Audiffred (1985: 174) and Neto et al. (2001: 403).

Corallina mediterranea Aresch = *Corallina officinalis* L. Audiffred and Weisscher (1984: 33). *Corallina officinalis*. Augier (1985: 100). SALVAGE ISLANDS: Price et al. (1986: 75) and John et al. (2004: 73). CANARY ISLANDS: Price et al. (1986: 75), Viera-Rodríguez (1987: 242), Viera-Rodríguez et al. (1987: 273), Afonso-Carrillo and Sansón (1999: 195), Gil-Rodríguez et al. (2003: 25) and John et al. (2004: 73). CAPE VERDE ISLANDS: Price et al. (1986: 75).

Haliptilon

The generic concept of this genus is not consensual. For the relationship between *Haliptilon* and other Janieae genera, see *Cheilosporum* entry. According to Womersley and Johansen (1996: 296), *Haliptilon* distinguishes from other Janieae genera by possessing a thallus pinnate with compressed axial intergenicula without lobes, usually bearing two or more subterete branchlets, sometimes adventitiously; conceptacles in terminal branchlets, usually single and with apical pores; spore-producing conceptacles with 1–3 branchlets.

Haliptilon cubensis (Montagne ex Kützing)
Garbary & H. W. Johansen

Basionym: *Jania cubensis* Montagne ex Kützing, 1849: 709–710. *Type:* Montagne Herbarium (Agardh 1852); illustrations: Garbary and Johansen (1982, Fig. 10, p. 216) as *Haliptilon cubensis* (Montagne ex Kützing) Garbary & H. W. Johansen. *Type locality:* Cuba (Kützing 1849: 710).

Current placement/name: *H. cubensis* (Montagne ex Kützing) Garbary & H. W. Johansen. Examination of a fragment of the type material revealed stable characters at the genus level (Garbary and Johansen 1982: 212), but the status as a distinct species within genus remains uncertain until comparative studies of relevant types are done. Following their suggestion to merge material of *Cheilosporum* and *Haliptilon* within the *Jania* genus (Kim et al. 2007, see *Cheilosporum* entry), these authors suggested the new combination *J. cubensis* (Montagne ex Kützing 1849, p. 709–10). According to Afonso-Carrillo (unpublished results), plants from the Canary Islands reported as *Corallina cubensis* (Montagne ex Kützing) Kützing or *H. cubensis* are not morphologically different from *Haliptilon virgatum* (Zanardini) Garbary & H. W. Johansen

Published records:

CANARY ISLANDS: *Corallina cubensis*. Delgado et al. (1984: 105) and Jorge et al. (1984: 118).

Corallina cubensis (Montagne) Kützing. Gil-Rodríguez and Afonso-Carrillo (1980: 36), Afonso-Carrillo et al. (1983: 30, 1984: 32) and Price et al. (1992: 125). *Haliptilon cubense* (Montagne ex Kützing) Garbary & H. W. Johansen. Gil-Rodríguez et al. (2003: 25) and John et al. (2004: 86).

CAPE VERDE ISLANDS: *Corallina cubensis* Montagne ex Kützing. Askenasy (1896: 26) and Price et al. (1992: 125).

Jania cubensis Montagne ex Kützing. Price et al. (1992: 125).

Haliptilon purpuratum (Lamarck) Garbary & H. W. Johansen

Basionym: *Corallina purpurata* Lamarck 1815: 237. *Type:* PC; illustrations: not found. *Type locality:* Tenerife, the Canaries (Lamarck 1815: 237).

Current placement/name: *H. purpuratum* (Lamarck) Garbary & H. W. Johansen. Examination of the thallus surface of a fragment of the type material revealed stable characters at the genus level (Garbary and Johansen 1982: 212), but the status as a distinct species within genus remains uncertain until comparative studies of relevant types are undertaken. Afonso-Carrillo and Sansón (1999: 153) mentioned that according to the original description of *C. purpurata* and the variable morphology of *Haliptilon virgatum* (Zanardini) Garbary & H. W. Johansen, these two species are not distinct and a revision of both species based on examination of type material is recommended.

Published records:

CANARY ISLANDS: *Haliptilon purpuratum* (Lamarck) Garbary & H. W. Johansen. Afonso-Carrillo and Sansón (1999: 153, 195).

Haliptilon squamatum (Linnaeus) Johansen, L. Irvine & Webster

Basionym: *Corallina squamata* Linnaeus 1758: 806. *Lectotype:* Ellis's illustrations designated by Irvine and Johansen (1994: 50); illustrations: Ellis (1755, pl. 24, Fig. c.C, p. 63). *Type locality:* European Seas (Lamouroux 1816: 287).

Current placement/name: *Haliptilon squamatum* (Linnaeus) Johansen, L. Irvine & Webster according to Johansen et al. (1973: 212) and Irvine and Johansen (1994: 49) based on the study of lectotype material. Prud'homme van Reine et al. (1994: 81) examining the material collected by Piccone from Azores (1889), the Canaries (1884) and Madeira (Grunow 1868) housed at PAD Herbarium considered it as *H. virgatum* (Zanardini) Garbary & H. W. Johansen,

following the concept of Afonso-Carrillo and Sansón (1989: 25). Nevertheless, the type material was not observed. The presence of *H. squamatum* in the Archipelagos of Madeira and the Canaries is dubious and needs confirmation (Afonso-Carrillo et al. 1983: 46; Neto et al. 2001: 412). Kim et al. (2007: 1317) based on molecular and reproductive features proposed the new combination *Jania squamata* (Linnaeus) J. H. Kim, Guiry & H.-G. Choi.

Published records:

AZORES: *Corallina squamata* J. Ellis & Solander. Schmidt (1929b: 171, 1931: 65) and Feldmann (1946: 419).

Haliptilon squamatum. Neto (1994: 26).

Haliptilon squamatum (Linnaeus) H.W. Johansen., L.M. Irvine & A.M. Webster. South and Tittley (1986: 42) and Tittley and Neto (1994: 9, 2005: 250).

MADEIRA: *Corallina squamata*. Grunow (1868: 78).

Corallina squamata J. Ellis & Solander. Grunow (1868: 78) and Schmidt (1929b: 171, 1931: 101).

Haliptilon squamatum (Linnaeus) H. W. Johansen, L.M. Irvine & A.M. Webster. John et al. (2004: 86).

CANARY ISLANDS: 1 *Haliptilon squamatum* (Linnaeus) H. W. Johansen, L. M. Irvine & A. M. Webster. Afonso-Carrillo et al. (1983: 46), Price et al. (1992: 125), Irvine and Johansen (1994: 49), Irvine and Johansen (1994: 49), Afonso-Carrillo and Sansón (1999: 195), Gil-Rodríguez et al. (2003: 25) and John et al. (2004: 86).

Haliptilon virgatum (Zanardini) Garbary & H. W. Johansen

Basionym: *Corallina virgata* Zanardini 1844: 1025. *Type:* MCVE (Stafleu and Cowan 1988: 516); illustrations: not found. *Type locality:* Adriatic Sea (Zanardini 1844: 1025).

Current placement/name: *H. virgatum* (Zanardini) Garbary & H. W. Johansen. Examination of a fragment of the type material revealed stable characters at the genus level (Garbary and Johansen 1982: 212), but the status as a distinct species within genus remains uncertain until comparative studies of relevant types are done. Kim et al. (2007) based on molecular and reproductive features proposed to merge *Haliptilon* within *Jania* genus (see *Cheilosporum* entry) but did not suggest a new combination for *H. virgatum*.

For the relationship between *H. virgatum* and *Corallina lobata* J. V. Lamouroux, see *C. lobata* entry.

Published records:

AZORES: Piccone (1889: 210) and Neto and Tittley (1995: 490).

Haliptilon virgatum (Zanardini) Garbary & H. W. Johansen. South and Tittley (1986: 42), Prud'homme van Reine et al. (1994: 81), Tittley and Neto (1994: 9), Tittley et al. (1998: 468) and Wallenstein and Neto (2006: 199).

MADEIRA: *Haliptilon virgatum* (Zanardini) Garbary & H. W. Johansen. Audiffred and Prud'homme van Reine (1985: 41) and John et al. (2004: 87).

SALVAGE ISLANDS: Price et al. (1992: 125).

CANARY ISLANDS: Piccone (1889: 210) and Price et al. (1992: 125).

Haliptilon virgata (Zanardini) Garbary & H. W. Johansen. Viera-Rodríguez and Wildpret de la Torre (1986: 217), Price et al. (1992: 125), Prud'homme van Reine et al. (1994: 94), Afonso-Carrillo and Sansón (1999: 195), Rojas-González and Afonso-Carrillo (2002b: 100), Gil-Rodríguez et al. (2003: 25) and Muñoz et al. (2007: 107).

Haliptilon virgatum. Gil-Rodríguez et al. (1992: 112) and Prud'homme van Reine et al. (1994: 81).

Haliptilon virgata. Afonso-Carrillo and Sansón (1989: 25).

Haliptilon virgatum (Zanardini) Garbary & H. W. Johansen. Elejabeitia et al. (1992: 8), Pinedo et al. (1992: 39), Guadalupe et al. (1995: 37), Haroun et al. (2003: 108), Montañés et al. (2003: 124), Sangil et al. (2003a: 103, b, 2004: 89, 311, 2005: 326) and John et al. (2004: 87).

Jania

The generic concept of this genus is not consensual. For the relationship between *Jania* and other Janieae genera, see *Cheilosporum* entry. According to Womersley and Johansen (1996: 296), *Jania* distinguishes from other Janieae genera by possessing a dichotomous thallus with terete or compressed intergenicula bearing axial conceptacles.

Jania adhaerens J. V. Lamouroux

Basionym: *Jania adhaerens* J. V. Lamouroux 1816: 270. *Holotype*: CN (Johansen 1971: 247; Cribb 1983); illustrations: not found. *Type locality*: Mediterranean (Lamouroux 1816: 270).

Current placement/name: *J. adhaerens* J. V. Lamouroux but status and disposition uncertain as the type was not studied in a modern context.

Published records:

AZORES: South and Tittley (1986: 42), Prud'homme van Reine (1988: 179), Neto (1994: 26, 2001: 104),

Tittley and Neto (1994: 9) and Tittley et al. (1998: 468).

MADEIRA: John et al. (2004: 93).

SALVAGE ISLANDS: Price et al. (1992: 135), Kristiansen et al. (1993: 95) and John et al. (2004: 93).

CANARY ISLANDS: Afonso-Carrillo (1980b: 55), Gil-Rodríguez and Afonso-Carrillo (1980: 36), Afonso-Carrillo et al. (1984: 30), González et al. (1986: 319), Afonso-Carrillo and Sansón (1989: 26, 1999: 196), Reyes and Sansón (1991: 79), Ballesteros et al. (1992: 520), Pinedo et al. (1992: 40), Price et al. (1992: 135), Ballesteros (1993: 21), Kristiansen et al. (1993: 95), Guadalupe et al. (1995: 37), Reyes et al. (2000: 140), Haroun et al. (2002: 146, 2003: 110), Rojas-González and Afonso-Carrillo (2002b: 102), Gil-Rodríguez et al. (2003: 25), Montañés et al. (2003: 124), Sangil et al. (2003a: 103, b: 311), John et al. (2004: 93), Sangil et al. (2004: 89, 2005: 326) and Muñoz et al. (2007: 107).

CAPE VERDE ISLANDS: Otero-Schmitt and Sanjuan (1992: 382), Price et al. (1992: 135) and Otero-Schmitt (1993: 47).

Jania capillacea Harvey

Basionym: *Jania capillacea* Harvey 1853: 84–85. *Holotype*: TCD (Toumey 70; Dawson 1953: 116; Cribb 1983); illustrations: not found. *Type locality*: Honda Bay, Florida, USA (Silva et al. 1996: 241).

Current placement/name: *J. capillacea* Harvey but status and disposition uncertain as the type was not studied in a modern context. Similarities between this species and *J. adhaerens* J. V. Lamouroux are discussed in Cribb (1983) and Schneider and Searles (1991). Further comments on the relation between *J. capillacea* and *Jania pumila* J. V. Lamouroux for the Canary Islands in *J. pumila* entry.

Published records:

AZORES: Neto (2001: 104) and Wellenstein and Neto (2006: 1999).

MADEIRA: John et al. (2004: 93).

SALVAGE ISLANDS: Price et al. (1992: 135) and John et al. (2004: 93).

CANARY ISLANDS: Afonso-Carrillo and Sansón (1989: 26, 1999: 196), Reyes and Sansón (1991: 79), Pinedo et al. (1992: 40), Price et al. (1992: 135), Ballesteros (1993: 22), Prud'homme van Reine et al. (1994: 94), Gil-Rodríguez et al. (2003: 25), Haroun et al. (2003: 110), Sangil et al. (2003a: 103, b: 311) and John et al. (2004: 93).

Jania capillacea J. V. Lamouroux. Afonso-Carrillo et al. (1983: 46, 1984: 30) and Price et al. (1992: 135).

CAPE VERDE ISLANDS: Otero-Schmitt and Sanjuan (1992: 382), Price et al. (1992: 135) and Otero-Schmitt (1993: 51).

Jania crassa J. V. Lamouroux

Basionym: *Jania crassa* J. V. Lamouroux 1821: 23. **Holotype:** CN Lamouroux Herbarium under *Jania* in folder c8-f74 (Woelkerling and Nelson 2004: 53); illustrations: Lamouroux (1821, pl. 69, Figs. 9–10, p. 23) and Johansen (1971, Fig. 10, p. 244) as *J. crassa*; Johansen and Womersley (1994, Fig. 37, p. 622) as *Jania verrucosa* J. V. Lamouroux; see comments. **Type locality:** Dusky Sound, South Island, New Zealand (Lamouroux 1821: 23).

Current placement/name: *J. verrucosa* J. V. Lamouroux but status and disposition uncertain as the type of *J. verrucosa* was not studied in a modern context. Johansen and Womersley (1994: 617) and Womersley and Johansen (1996: 305, 307) considered *J. crassa* as a heterotypic synonym of *J. verrucosa* stating that they are almost “certainly conspecific at least in southern Australia”. This conclusion was only based on the study of the holotype of *J. crassa*. However, according to Woelkerling and Nelson (2004: 65), the proposed conspecificity needs to be confirmed through a comparative study of both types.

Published records:

AZORES: South and Tittley (1986: 43), Neto (1994: 26) and Tittley and Neto (1994: 9).

Jania crassa. Prud'homme van Reine (1988: 179), Stengenga et al. (1997: 585) and Neto (2000a: 140, b: 490).

CAPE VERDE ISLANDS: Otero-Schmitt and Sanjuan (1992: 384) and Price et al. (1992: 136).

Jania intermedia (Kützing) P. C. Silva

Basionym: *Corallina intermedia* Kützing 1858: 37–38, 42. **Type:** not found; illustrations: Kützing (1858, pl. 79, Fig. I, p. 36–38; pl. 86, Fig. IV, p. 42). **Type locality:** Cape of Good Hope, South Africa (Silva et al. 1996: 242).

Current placement/name: *J. intermedia* (Kützing) P. C. Silva but status and disposition uncertain as the type was not studied in a modern context. Silva et al. (1996: 242) mentioned that the correct combination is *J. intermedia* and not *Jania intermedia* Kützing which was only a bibliographic reference and not the name of the basionym. Neto et al. (2001: 412.) studying Madeira specimens cited it as doubtful record mentioning that “it might be a variety of *J. rubens* (Linnaeus) J.V. Lamouroux”. According to Woelkerling and Nelson (2004: 57), this name should be considered a *nomen nudum* as there are no valid diagnosis descriptions.

Published records:

MADEIRA: *Corallina intermedia* Grunow. Neto et al. (2001: 412).

Jania longifurca Zanardini

Basionym: *Jania longifurca* Zanardini 1844: 1025. **Type:** probably in MCVE Zanardini's collections; illustrations: not found. **Type locality:** Dalmatia, Italy (Zanardini 1844: 1025).

Current placement/name: *J. longifurca* Zanardini but status and disposition uncertain as the type was not studied in a modern context. Sansón et al. (1991: 531) mentioned that this species is poorly known in the Canary Islands and needs further studies. The designation/localization of the type material and its study in a modern context is required.

Published records:

AZORES: Feldmann (1946: 421), South and Tittley (1986: 43), Neto (1994: 26), Tittley et al. (1998: 468) and Neto (2000a: 140, b: 490, 2001: 105).

MADEIRA: Levring (1974: 70), Sansón et al. (1991: 530) and Neto et al. (2001: 404).

CANARY ISLANDS: Sansón et al. (1991: 530), Afonso-Carrillo and Sansón (1999: 196), Haroun et al. (2002: 146) and Gil-Rodríguez et al. (2003: 25).

Jania micrarthrodia J. V. Lamouroux

Basionym: *Jania micrarthrodia* J. V. Lamouroux 1816: 271. **Holotype:** CN Lamouroux Herbarium; illustrations: Lamouroux (1816, pl. 9, Fig. 5a; pl. 69, Figs. 7–8, p. 271); Johansen and Womersley (1994, Figs. 32–36, p. 622). **Type locality:** Australasia (Lamouroux 1816: 271).

Current placement/name: *J. micrarthrodia* J. V. Lamouroux according to Johansen and Womersley (1994: 611–613) based on the study of type material. Gil-Rodríguez et al. (2003: 212) mentioned that the material of *J. micrarthrodia* for the Canaries could not be located.

Published records:

CANARY ISLANDS: Price et al. (1992: 136) and Afonso-Carrillo and Sansón (1999: 196).

Jania natalensis Harvey

Basionym: *Jania natalensis* Harvey 1849: 107. **Type:** TCD (Johansen 1971: 247); illustrations: not found. **Type locality:** Port Natal, South Africa (Johansen 1971: 247).

Current placement/name: *Jania verrucosa* J. V. Lamouroux but status and disposition uncertain as the type was not studied in a modern context. Johansen and Womersley

(1994: 616) merged *J. natalensis* in synonymy with *J. verrucosa* but no revision of the type material was made. Agardh (1870: 363) cited *J. natalensis* as doubtful to the Azores (Santa Maria).

Published records:

- AZORES: Schmidt (1929b: 172, 1931: 66).
- Corallina natalensis* Kützing. Neto (2000b: 485).
- Jania natalensis*? Agardh (1870: 363) and Piccone (1889: 214).
- Jania? Natalensis* Harvey. Gain (1914: 22).

Jania pumila J. V. Lamouroux

Basionym: *Jania pumila* J. V. Lamouroux 1816: 269. *Holotype*: CN Lamouroux Herbarium; illustrations: Lamouroux (1816, pl. 9, Fig. 2, p. 269). *Type locality*: Red Sea and Western Indies (Lamouroux 1816: 269).

Current placement/name: *J. pumila* J. V. Lamouroux but status and disposition uncertain as the type was not studied in a modern context. The holotype specimen of *J. pumila* housed in Lamouroux's Herbarium in Caen is in depauperate and/or sterile conditions being difficult to interpret (Johansen and Womersley 1994: 610; Abbott 1999: 189). Villena-Balsa et al. (1987: 20) revising material from the Canary Islands identified it provisionally as *J. pumila*. They mentioned that the material could belong to other species also characterized by reduced-size plants and not correctly described e.g. *Jania gibbosa* J. V. Lamouroux, *J. pygmea* J. V. Lamouroux, *J. compressa* J. V. Lamouroux and *J. capillacea* Harvey.

Published records:

- MADEIRA: Levring (1974: 70), González (1978: 48) and Neto et al. (2001: 404).
- SALVAGE ISLANDS: Price et al. (1992: 136) and John et al. (2004: 93).
- CANARY ISLANDS: Price et al. (1992: 136), Afonso-Carrillo et al. (1978: 232, 1983: 30, 1984: 30), González (1978: 48), Gil-Rodríguez and Afonso-Carrillo (1980: 36), Pinedo et al. (1992: 40), Afonso-Carrillo and Sansón (1999: 196), Haroun et al. (2003: 112), Gil-Rodríguez et al. (2003: 25), Sangil et al. (2003a: 103, 2005: 326) and John et al. (2004: 93).
- Jania pumila*. Afonso-Carrillo and Sansón (1989: 25) and Price et al. (1992: 136).

Jania rubens (Linnaeus) J. V. Lamouroux var. *corniculata* (Linnaeus) Yendo

Basionym: *Corallina corniculata* Linnaeus 1758: 806. *Lectotype*: LINN 1293.19 (Irvine and Johansen 1994: 56);

illustrations: not found. *Type locality*: European Ocean (Irvine and Johansen 1994: 56).

Current placement/name: *Jania rubens* (Linnaeus) J. V. Lamouroux var. *corniculata* (Linnaeus) Yendo but status and disposition uncertain as the type was not studied in a modern context. Irvine and Johansen (1994: 56) in their revision of the British Isles material considered the name *J. rubens* var. *corniculata* (Linnaeus) Yendo as valid but without checking the type material. Recently, John et al. (2004: 93) considered *J. corniculata* (Linnaeus) J. V. Lamouroux as a synonym of *Jania rubens* (Linnaeus) J. V. Lamouroux but did not provide any taxonomic comments. In their integrative taxonomic approach including phylogeny reconstructions based on the cox1 and 18S rRNA genes in addition to a comparative morphological assessment study Walker et al. (2009) observed that *J. rubens* var. *rubens* and *J. rubens* var. *corniculata* clustered together in both phylogenies suggesting that for those genes there was no genetic basis for the morphological variation. These conclusions however were not based on type material observations.

Published records:

- AZORES: Gain (1914: 22), Tittley et al. (1998: 468) and Neto (2000b: 485, 2001: 105).
- Jania corniculata* J. V. Lamouroux. Piccone (1889: 213).
- Jania corniculata* (Linnaeus) J. V. Lamouroux. South and Tittley (1986: 42).
- Jania cf. corniculata* (Linnaeus) J.V. Lamouroux. Neto (1994: 26) and Tittley and Neto (1994: 9).
- Jania rubens* J. V. Lamouroux var. *corniculata* (Linnaeus) Yendo. Schmidt (1931: 66).

MADEIRA: *Jania cf. corniculata*. Neto et al. (2001: 404). *Jania corniculata* (Linnaeus) J. V. Lamouroux. Audiffred and Prud'homme van Reine (1985: 43).

SALVAGE ISLANDS: *Jania cf. corniculata*. John et al. (2004: 93).

CANARY ISLANDS: *Jania corniculata* (Linnaeus) J. V. Lamouroux. Gil-Rodríguez and Afonso-Carrillo (1980: 36), Afonso-Carrillo et al. (1983: 30, 1984: 30) and Price et al. (1992: 135).

Jania corniculata. Afonso-Carrillo and Sansón (1989: 26). *Jania rubens* (Linnaeus) J. V. Lamouroux var. *corniculata* (Linnaeus) Yendo. Price et al. (1992: 135) and Irvine and Johansen (1994: 57).

Jania rubens (Linnaeus) J. V. Lamouroux var. *rubens*

Basionym: *Corallina rubens* Linnaeus 1758: 806. *Lectotype*: UPS Herb. Burser vol. XX, 72; designated by Irvine and Johansen (1994: 56); illustrations: Lamouroux (1816, pl.

9, figs 6–7, p. 272) as *Jania rubens* (Linnaeus) J. V. Lamouroux. *Type locality*: European Ocean (Linnaeus 1758: 806).

Current placement/name: *J. rubens* (Linnaeus) J. V. Lamouroux var. *rubens*, a conclusion of Irvine and Johansen (1994: 56) after studying the lectotype in a modern context. Price et al. (1992: 136) suggested that *J. rubens* and *Corallina rubens* J. Ellis & Solander in González (1977a: 26) are synonyms. Prud'homme van Reine et al. (1994: 94) confirmed the diagnosis of the *J. rubens* material from Cape Verde Islands collected by Piccone (1886) and housed at PAD and W Herbarium but recognized that material was from Tenerife (the Canaries) and not from Cape Verde Islands. They followed Taylor's (1960) concept and did not review the type material.

Published records:

AZORES: Gain (1914: 15, 22) and Prud'homme van Reine et al. (1994: 94).

Jania rubens. Schmidt (1929a: 331).

Jania rubens (Linnaeus) J. V. Lamouroux. Feldmann (1946: 417), Castro and Viegas (1983: 20, 1987: 63), Weisscher (1983: 70), South and Tittley (1986: 43), Tittley and Neto (1994: 9), Neto (2000a: 140, 2000b: 490–491), Toste et al. (2003: 1270) and Wallenstein and Neto (2006: 199).

Jania rubens J. V. Lamouroux. Schmidt (1931: 66).

Jania rubens. Agardh (1870: 363), Palminha (1957: 66), Neto (1994: 26, 2000b: 490) and Neto and Tittley (1995: 489).

Jania rubens (Linnaeus) Areschoug. Lemoine (1924: 130).

Jania rubens (Linnaeus) J. V. Lamouroux var. *rubens*. Audiffred and Prud'homme van Reine (1985: 43) and Irvine and Johansen (1994: 55).

MADEIRA: *Corallina rubens* J. V. Lamouroux. Gain (1914: 10) and John et al. (2004: 93).

Jania rubens J. V. Lamouroux. Piccone (1884: 10), Gain (1914: 9) and Schmidt (1931: 101).

Jania rubens (Linnaeus) J. V. Lamouroux. Grunow (1868: 78) and Levring (1974: 70).

Jania rubens. Johnston (1969: 215), Augier (1985: 100) and Afonso-Carrillo and Sansón (1989: 26).

Jania rubens (Linnaeus) J. E. Areschoug. Lemoine (1924: 130).

SALVAGE ISLANDS: Price et al. (1992: 136) and John et al. (2004: 93).

CANARY ISLANDS: *Corallina (Jania) rubens* Linnaeus. Askenasy (1896: 26) and Prud'homme van Reine et al. (1994: 94).

Jania rubens. Johnston (1969: 215), González (1977b: 100), Delgado et al. (1984: 106), Gil-Rodríguez et al. (1992: 109) and Price et al. (1992: 136).

Jania rubens (Linnaeus) J. V. Lamouroux. Ardré (1970: 229), Santos Guerra et al. (1970: 24), Cordeiro-Marino (1978: 50), González (1978: 48), Afonso-Carrillo et al. (1978: 232, 1984: 30), Afonso-Carrillo and Gil-Rodríguez (1980: 165), Gil-Rodríguez and Afonso-Carrillo (1980: 36), López Hernández and Gil-Rodríguez (1981: 156), Weisscher (1983: 70), González et al. (1986: 319), Viera-Rodríguez and Wildpret de la Torre (1986: 216), Reyes and Sansón (1991: 79), Pinedo et al. (1992: 40), Medina and Haroun (1993: 110), Kristiansen et al. (1993: 95), Guadalupe et al. (1995: 37), Afonso-Carrillo and Sansón (1999: 196), Gil-Rodríguez et al. (2003: 25), Rojas-González and Afonso-Carrillo (2002a: 33), Haroun et al. (2003: 112), Montañés et al. (2003: 124), Sangil et al. (2003a: 103), John et al. (2004: 93), Rojas-González and Afonso-Carrillo (2004: 138) and Muñoz et al. (2007: 107).

Jania rubens J. V. Lamouroux. Vickers (1896: 306), Schmidt (1931: 101 and Price et al. (1992: 136).

Jania rubens (Linnaeus). Piccone (1884: 42, 56) and Acuña González (1972: 5).

Jania rubens. Price et al. (1992: 136).

Jania rubens (Linnaeus) J. V. Lamouroux var. *rubens*. Irvine and Johansen (1994: 55).

CAPE VERDE ISLANDS: *Corallina (Jania) rubens* Linneaus. Price et al. (1992: 136).

Jania rubens (Linnaeus) J. V. Lamouroux. Piccone (1886: 66), Price et al. (1992: 136) and Prud'homme van Reine et al. (1994: 94).

Jania rubens J. V. Lamouroux. Schmidt (1931: 101).

Jania cf. rubens (Linnaeus) J. V. Lamouroux. Otero-Schmitt and Sanjuan (1992: 382).

Jania tenella (Kützing) Grunow

Basionym: *Corallina tenella* Kützing 1858: 41. *Holotype*: L (Dawson 1953: 121); illustrations: Kützing (1858, pl. 85, fig. II, p. 41). *Type locality*: Naples, Italy (Silva et al. 1996: 244).

Current placement/name: *J. tenella* (Kützing) Grunow but status and disposition uncertain as the type was not studied in a modern context.

Published records:

CAPE VERDE ISLANDS: *Corallina (Jania) tenella* Kützing. Price et al. (1992: 136).

Jania verrucosa J. V. Lamouroux

Basionym: *Jania verrucosa* J. V. Lamouroux 1816: 270. **Holotype:** CN Lamouroux Herbarium (Womersley and Johansen 1996: 305); illustrations: Lamouroux (1816, pl. 9, Fig. 4 a.B, p. 270). **Type locality:** North America (Lamouroux 1816: 270).

Current placement/name: *J. verrucosa* J. V. Lamouroux but status and disposition uncertain as the type was not studied in a modern context. For the relationship between *J. verrucosa*, *J. natalensis* and *J. cryptarthrodia*, see *J. natalensis* and *J. cryptarthrodia* entries.

Published records:

AZORES: *Jania verrucosa*. Neto (2000b: 491, 2001:105).

CAPE VERDE ISLANDS: Price et al. (1992: 136).

Discussion

From the 31 names of species and infraspecific taxa of geniculate Corallinales recorded from the Macaronesian region, four are based on type material collected from localities within this region, while the rest (87%) are based on types from localities elsewhere (Table 1). All species are validly published (ICBN, Art. 6.2, 29.1, 42.3 and 60). The correct application of these names to species and specific taxa, however, is linked to an understanding of the type material and to the provisions of the ICBN (McNeill et al. 2006).

For *Haliptilon cubensis*, *H. virgatum* and the four names based on Macaronesian collections (*Corallina lobata*, *C. microptera*, *C. millegrana* and *Haliptilon purpurata*), only a fragment of the holotype material was studied. The generic placement of the taxa was confirmed by ultrastructural investigations but the status of the species could not be determined (Garbarek and Johansen 1982). As a result, published information does not include detailed specific morphological and anatomical accounts, and its nomenclature lacks the foundations essential for stability. Therefore, although those taxa are currently treated as distinct species by various authors, this treatment requires substantiation through morphological, anatomical and ultrastructural detailed studies of relevant type material.

From the list compiled in the present work, 15 taxa have not been studied in a modern context. *Jania intermedia* is a *nomen nudum* (ICBN, Art. 1) and therefore nomenclaturally illegitimate. The remaining 14 taxa, although listed as distinct species in various publications, require re-assessment through further studies of relevant type material. The same situation happens with the proposed heterotypic

synonyms. An example is the synonymy of *Corallina mediterranea* with *C. elongata* published by Irvine and Johansen (1994: 41). The segregating characters defined by these authors for *C. elongata*, related to reproductive and vegetative morphology, cannot be contrasted with the available original description of *C. mediterranea* from Areschoug (1852: 568–569). Also, the presence in the main axes of intergenicula as long as wide, described by Walker et al. (2009: 295–296) as distinguishing feature for *C. elongata*, does not agree with the description of *C. mediterranea*. The occurrence of most of these taxa in Macaronesia has yet to be verified as most records represent occasional findings from one single location.

Only ten types of Corallinales occurring in Macaronesia have been studied in a modern context. From these, two (*Jania crassa* and *Amphiroa verruculosa*) are considered heterotypic synonyms of other taxa. The heterotypic synonymy of *Jania crassa* with *Jania verrucosa* (Johansen and Womersley 1994), however, requires further studies comparing both types. The remaining eight species are considered to be distinct taxa. Their diagnosing characters were emended based on, respectively, vegetative and reproductive anatomy in the case of *Amphiroa* (Norris and Johansen 1981: 5–6), vegetative and reproductive morphology in *Corallina* (Walker et al. 2009: 295; Table 4, p. 296) and morphological and anatomical vegetative characters in *Jania* (Johansen and Womersley 1994: 606). The confirmation of these species in the Macaronesia requires the study of their voucher specimens in light of these new concepts.

To summarize, only 8 of the 31 species and infraspecific taxa of geniculate Corallinales listed for Macaronesia have been confirmed to represent distinct species. Three others have been found to be heterotypic synonyms of other taxa and one (*Jania intermedia*) is nomenclaturally illegitimate. There remains 19 taxa that have yet to be fully reassessed through detailed studies encompassing specific morphological, anatomical and ultrastructural investigations, preferably complemented with molecular studies. This task has many difficulties, starting with the fact that for three species (*Corallina deshayesii*, *C. granifera* and *C. intermedia*) the type collection whereabouts are uncertain. However, this kind of taxonomic work may confirm the molecular conclusions of Kim et al. (2007) suggesting that the genera *Cheilosporum*, *Haliptilon* and *Jania* (tribe Janieae) should be merged into a single genus *Jania* recognized by the generic characters listed by Johansen and Silva (1978: 413, Table 1, p. 414).

Conclusion

The present catalog and its analysis represent initial steps in creating a reliable database on the species diversity of

geniculate Corallinales in the Macaronesian region. Although 31 names of geniculate Corallinales have been recorded for this region, the actual number of species and genera present is uncertain and will continue to remain so until detailed monographic studies are undertaken. How many of the names previously used for the Macaronesian region taxa can correctly be applied to specimens for this region also remains uncertain as does the number of species present, but so far undetected. An accurate figure of the actual diversity for the Macaronesian geniculate corallines requires extensive work involving new collections, the detailed study of fresh material, historical collections and type material, complemented, if possible, with molecular evaluations.

Acknowledgments We thank Drs. Willem F. Prud'homme van Reine (National Herbarium Nederland-L) and Bruno de Reviers (Muséum National d'Histoire Naturelle-PC) who provided specialized old literature and/or type material information, Ian Tittley (Natural History Museum), José M. N. Azevedo and Gustavo Martins for the English revision of the manuscript, Nuno Álvaro for the study area map, and to Dr. Inka Bartsch and an anonymous reviewer for their useful comments and suggestions. This work was partly supported by the Programme AlBan, the European Union Programme of High Level Scholarships for Latin America (through scholarship E05D060221MX) and "Consejo Nacional de Ciencia y Tecnología" doctoral scholarship 176162, the UNAMUNO Programme of PhD Scholarships for Europe, and "Centro de Investigação de Recursos Naturais", University of the Azores.

References

- Abbott IA (1999) Marine red algae of the Hawaiian Islands. Bishop Museum Press, Honolulu
- Acuña González A (1972) Observaciones ecológicas sobre las algas de la zona litoral de Las Galletas Tenerife. Vieraea 1:2–5
- Afonso-Carrillo J (1980a) Algunas observaciones sobre la distribución vertical de las algas en la isla del Hierro (Canarias). Vieraea 10:3–16
- Afonso-Carrillo J (1980b) Nota sobre algunas Corallinaceae (Rhodophyta) nuevas para la flora fitológica de las Islas Canarias. Vieraea 10:53–58
- Afonso-Carrillo J (1986) Observaciones en *Amphiroa fragilissima* (L.) Lamouroux (Corallinaceae Rhodophyta) con el microscopio electrónico de barrido. Vieraea 16:189–192
- Afonso-Carrillo J (1988) Structure and reproduction of *Spongites wildpretii* sp. nov. (Corallinaceae, Rhodophyta) from the Canary Islands, with observations and comments on *Spongites absimile* comb. nov. Br Phycol J 23:89–102
- Afonso-Carrillo J, Gil-Rodríguez MC (1980) Datos para la flora marina de la isla de Fuerteventura. Vieraea 10:147–170
- Afonso-Carrillo J, Sansón M (1989) Clave ilustrada para la determinación de los macrófitos marinos bentónicos de las Islas Canarias. Departamento de Biología Vegetal (Botánica), Universidad de La Laguna, La Laguna
- Afonso-Carrillo J, Sansón M (1999) Algas hongos y fanerógamas marinas de las Islas Canarias. Clave analítica. Materiales Didácticos Universitarios. Serie Biología/2. Universidad de La Laguna, La Laguna
- Afonso-Carrillo J, Sobrino C (2004) Two amphi-Atlantic species of *Botryocladia* (Rhodymiales Rhodophyta) in the Canary Islands (Eastern Atlantic). Cryptogamie Algol 25:147–159
- Afonso-Carrillo J, Gil-Rodríguez MC, Wildpret de la Torre W (1978) Estudio de la vegetación algal de la costa del futuro polígono industrial de Granadilla (Tenerife). Vieraea 8:201–242
- Afonso-Carrillo J, Gil-Rodríguez MC, Haroun-Tabraue R, Villena-Balsa M, Wildpret de la Torre W (1983) Adiciones y correcciones al catálogo de algas marinas bentónicas para el Archipiélago Canario. Vieraea 13:27–49
- Afonso-Carrillo J, Gil-Rodríguez MC, Wildpret de la Torre W (1984) Algunas consideraciones florísticas corológicas y ecológicas sobre las algas Corallinaceae (Rhodophyta) de las islas Canarias. Anales Biol 2:23–37
- Afonso-Carrillo J, Sangil C, Sansón M (2009) *Lomentaria benahoarensis* (Lomentariaceae, Rhodophyta), a diminutive epiphytic new species from La Palma, Canary Islands (eastern Atlantic Ocean). Bot Mar 52:236–247
- Agardh JG (1852) Species generae et ordines algarum, seu descriptiones succinctae specierum, generum et ordinum, quibus algarum regnum constituitur. Volumen secundum: algas florideas complectens. Part 3, fasc. 1. C.W.K. Gleerup, Lundae [Lund]
- Agardh JG (1870) Om de under Korvetten Josephines expedition sistliden sommar insamlade Algerne. Ofver K Vet Ak Förhandl 27:359–366
- Ardisson F (1883) Phycologia mediterranea. Parte Prima Floridee. Mem Soc Cittog Ital 1:1–516
- Ardré F (1970) Contribution à l'étude des algues marines du Portugal I-La Flore. Port Acta Biol, ser B10:137–555
- Ardré F, Boudouresque CF, Cabioch J (1974) *Sympyocladia marchantiooides* (Harv.) Falkenberg (Rhodomélacées Ceramiales) aux Açores. Bull Soc phycol Fr 19:178–182
- Areschoug JE (1852) Ordo XII. Corallineae. In: Agardh JG (ed) Species generae et ordines algarum. Volumen secundum: algas florideas complectens. Lund, Sweden, pp 506–576
- Askenasy ME (1896) Énumeration des algues des îles du Cap Vert. Bol Soc Broteriana 13:1–26
- Audiffred PAJ (1985) Marine algae of El Hierro (Canary Islands). Vieraea 14:157–183
- Audiffred PAJ, Prud'homme van Reine WF (1985) Marine algae of Ilha do Porto Santo and Deserta Grande (Madeira Archipelago) (Cancap Project Contributions No. 40). Bolm Mus munic Funchal 37:20–51
- Audiffred PAJ, Weisscher FLM (1984) Marine algae of Salvagem Grande (Salvage Island Macaronesia) (Cancap Project Contribution No. 37). Bolm Mus munic Funchal 36:5–37
- Augier H (1985) Première contribution à la cartographie des biocénoses marines benthiques de l'île de Madère. Bolm Mus munic Funchal 37:86–129
- Bailey JC (1999) Phylogenetic positions of *Lithophyllum incrustans* and *Titanoderma pustulatum* (Corallinaceae, Rhodophyta) based on 18S rRNA gene sequence analyses, with a revised classification of the Lithophylloideae. Phycologia 38:208–216
- Ballesteros E (1993) Algunas observaciones sobre las comunidades de algas profundas en Lanzarote y Fuerteventura (Islas Canarias). Vieraea 22:17–27
- Ballesteros E, Sansón M, Reyes J, Afonso-Carrillo J, Gil-Rodríguez MC (1992) New records of benthic marine algae from the Canary Islands. Bot Mar 35:513–522
- Barton ES (1897) Welwitsch's African marine algae. J Bot, Lond 35:369–374
- Børgesen F (1929) Marine algae from the Canary Islands especially from Teneriffe and Gran Canaria II. Rhodophyceae. Part II. Cryptonemiales, Gigartinales and Rhodynemiales. Biol Meddr K danske Vidensk Skr 8:1–97
- Bornet E (1892) Les algues de P.-K.-A. Schousboe recoltées au Maroc et dans la Méditerranée de 1815 à 1829. Mém Soc natn Math Cherbourg 28:165–376
- Castro ML, Viegas MC (1983) Estudo dos povoamentos de algas fotófilas da Ilha de S. Miguel (Açores). 1. Resultados preliminares

- sobre a facies de *Corallina elongata* Ellis and Solander. Arquipélago 4:7–30
- Castro ML, Viegas MC (1987) Contribuição para o estudo da zona intertidal (substrato rochoso) da ilha de São Miguel-Acores. Fácies de *Corallina elongata* Ellis and Solander. Resultados preliminares. Cuad Marisq Publ 11:59–69
- Chapman G (1955) Aspects of the fauna and flora of the Azores. VI. Density of animal life in the coralline alga zone. Ann Mag Nat Hist 12:801–805
- Cordeiro-Marino M (1978) Rodofíceas bentônicas marinhas do Estado de Santa Catarina. Rickia 7:1–243
- Cremades J, Bárbara I, Veiga JA (1997) *Amphiroa van-bosseae* (Corallinales Rhodophyta) on European Atlantic coasts. Cryptogamie Algol 18:11–17
- Cribb AB (1983) Marine algae of the southern Great Barrier Reef Part. 1. Rhodophyta. Australian coral reef Society, Brisbane
- Dawson EY (1953) Marine red algae of Pacific Mexico. Part 1. Bangiales to Corallinaceae subf. Corallinoidae. Allan Hancock Pacif Exped 17:1–239
- Delgado E, González MN, Jorge D (1984) Contribución al estudio de la vegetación ficológica de la zona de Arinaga (Gran Canaria). Botanica Macaron 12–13:97–110
- Elejabeitia Y, Reyes J, Afonso-Carrillo J (1992) Algas marinas bentônicas de Punta del Hidalgo, Tenerife (Islas Canarias). Vieraea 21:1–28
- Ellis J (1755) An essay towards a natural history of the corallines and other marine productions of the like kind commonly found on the coasts of Great Britain and Ireland. To which is added the description of a large marine polype taken near the North Pole by the Whale-fishers in the summer 1753. Privately Published, London
- Ellis J, Solander D (1786) The natural history of many curious and uncommon zoophytes collected from various parts of the globe by the late John Ellis. Systematically arranged and described by the late Daniel Solander. B. White and Son, London
- Feldmann J (1946) La flore marine des îles Atlantides. Mém Soc Biogéogr 8:395–435
- Gain L (1914) Algues provenant des campagnes de "l'Hirondelle" II' (1911–1912). Bull Inst Océanogr Monaco 279:1–23
- Garbarry DJ, Johansen HM (1982) Scanning electron microscopy of *Corallina* and *Haliptilon* (Corallinaceae Rhodophyta): surfaces features and their taxonomic implications. J Phycol 18:211–219
- Gil-Rodríguez MC (1980) Revisión taxonómica-ecológica del género *Cystoseira* C. Ag. en el Archipiélago Canario. Vieraea 9:115–148
- Gil-Rodríguez MC, Afonso-Carrillo J (1980) Catálogo de las algas marinas bentônicas (Cyanophyta, Chlorophyta, Phaeophyta y Rhodophyta) para el Archipiélago Canario (Tenerife). Aula de Cultura, Excmo. Cabildo Insular, Santa Cruz de Tenerife
- Gil-Rodríguez MC, Afonso-Carrillo J, Wildpret de la Torre W (1987) Praderas marinas de *Zostera noltii* (Zosteraceae) en las Islas Canarias. Vieraea 17:43–416
- Gil-Rodríguez MC, Afonso-Carrillo J, Haroun RJ (1992) Flora ficológica de las islas Canarias. In: Kunkel G (ed) Flora y vegetación del Archipiélago Canario. Tratado Florístico 1ª Parte. EDIRCA, Las Palmas de Gran Canaria, pp 95–121
- Gil-Rodríguez MC, Haroun RJ, Ojeda RA, Berecibar ZE, Domínguez SP, Herrera MB (2003) Protocista. In: Moro L, Martín JL, Garrido MJ, Izquierdo I (eds) Lista de especies marinas de Canarias (algas, hongos, plantas y animales). Consejería de Política Territorial y Medio Ambiente del Gobierno de Canarias, pp 17–33
- González N (1977a) Estudio de la vegetación litoral de la zona de Mas-palomas. Botanica Macaron 4:23–30
- González N (1977b) Estudio de la vegetación bentónica litoral del nordeste de la isla de Gran Canaria (Bañaderos, San Felipe, Sardina, Las Nieves). Botanica Macaron 4:85–104
- González N (1978) Estudio algológico de la Playa del Burrero (Gran Canaria). Botanica Macaron 6:43–51
- González N, Rodrigo PJD, Suárez RC (1986) Flora y vegetación del Archipiélago Canario. Las Palmas de Gran Canaria, MAE/Spain
- Grunow A (1868) Algen. In: Fenzl E (ed) Reise der Österreichischen Fregatte Novara um die Erde in den Jahren 1857 1858 und 1859 unter den Befehlen des Commodore B. von Wüllerstorff-Urbair. Botanischer Theil 1. Sporenplatten. Kaiserl. -Königl. Hof- und Staatsdruckerei, Wien, pp 1–104
- Guadalupe ME, Gil-Rodríguez MC, Hernández-González MC (1995) Fitobentos de Arrecife de Lanzarote, Reserva de la Biosfera (Islas Canarias). Cryptogamie Algol 16:33–46
- Hamel G, Lemoine P (1953) Corallinacées de France et d'Afrique du Nord. Archs Mus Hist nat Paris 7:1:15–136
- Hardisson A, Frias I, De Bonis A (1998) Mercury in algae of the Canary Islands littoral. Enviro Inter 24:945–950
- Haroun RJ, Gil-Rodríguez MC, Afonso-Carrillo J, Wildpret de la Torre W (1984) Vegetación bentónica del Roque de Los Órganos (Gomera). Anales Biol 2:107–117
- Haroun RJ, Gil-Rodríguez MC, Díaz de Castro J, Prud'homme van Reine WF (2002) A checklist of the marine plants from the Canary Islands (central eastern Atlantic Ocean). Bot Mar 45:139–169
- Haroun RJ, Gil-Rodríguez MC, Wildpret de la Torre W (2003) Plantas marinas de las Islas Canarias. Canseco editores, Talavera de la Reina
- Harvey WH (1849) Nereis australis or algae of the southern ocean: being figures and descriptions of marine plants collected on the shores of the Cape of Good Hope the extra-tropical Australian colonies Tasmania New Zealand and the Antarctic regions; deposited in the Herbarium of the Dublin University. Part 2. Reeve Brothers, London
- Harvey WH (1853) Nereis boreali-americana; or, contributions towards a history of the marine algae of the atlantic and pacific coasts of North America. Part II. Rhodospermeae. Smithson Contr Knowl 5:[i–ii], [1]–258
- Harvey AS, Broadwater ST, Woelkerling WJ, Mitrovski PJ (2003) *Choreonema* (Corallinales, Rhodophyta): 18S rDNA phylogeny and resurrection of the Hapalidiaceae for the subfamilies Choreonematoideae, Austrolithoideae and Melobesioidae. J Phycol 39:988–998
- Harvey AS, Woelkerling WJ, Millar AJK (2009) The genus *Amphiroa* (Lithophylloideae, Corallinaceae, Rhodophyta) from the temperate coasts of the Australian continent, including the newly described *A. klochkovana*. Phycologia 48:258–290
- Hebert PDN, Cywinski A, Ball SL, DeWaard JR (2003) Biological identifications through DNA barcodes. Proc Biol Sci, Royal Soc (B) 270:313–321
- Holmgren PK, Holmgren NH (1998) [continuously updated]. Index herbariorum: a global directory of public herbaria and associated staff. New York Botanical Garden's Virtual Herbarium. Available at: <http://sweetgum.nybg.org/ih/>
- Irvine LM, Johansen HW (1994) Corallinoideae. In: Irvine LM, Chamberlain YM (eds) Seaweeds of the British Isles, vol 1 Rhodophyta. Part 2B Corallinales Hildenbrandiales. HMSO Books, The Natural History Museum, London, pp 37–57
- Johansen HW (1971) Changes and additions to the articulated coralline flora of California. Phycologia 10:241–249
- Johansen HW (1981) Coralline Algae, a first synthesis. CRC Press, Boca Raton
- Johansen HW, Silva PC (1978) Janiaeae and Lithotrichaeae: two new tribes of articulated Corallinaceae (Rhodophyta). Phycologia 17:413–417
- Johansen HW, Womersley HBS (1994) *Jania* (Corallinales Rhodophyta) in southern Australia. Aust Syst Bot 7:605–625
- Johansen HW, Irvine LM, Webster AM (1973) *Haliptilon squamatum* (L.) comb. nov., a poorly known British coralline alga. Br Phycol J 8:212

- John DM, Prud'homme van Reine WF, Lawson GW, Kostermans TB, Price JH (2004) A taxonomic and geographical catalogue of the seaweeds of the western coast of Africa and adjacent islands. *Beih Nova Hedwigia* 127:1–339
- Johnston CS (1969) Studies on the ecology and primary production of Canary Islands marine algae. *Proc Int Seaweed Symp* 6:213–222
- Jorge D, González MN, Delgado E (1984) Macrofitobentos del litoral del Puerto de Las Nieves (Gran Canaria). *Botanica Macaron* 12–13:111–122
- Kim JH, Guiry MD, Oak JH, Choi DS, Kang SH, Chung H, Choi HG (2007) Phylogenetic relationships within the tribe Janieae (Corallinales, Rhodophyta) based on molecular and morphological data: a reappraisal of *Jania*. *J Phycol* 43:1310–1319
- Kohlmeyer J (1967) Intertidal and phycophilous fungi from Tenerife (Canary Islands). *Trans Br Mycol Soc* 50:137–147
- Kristiansen A, Nielsen R, Pedersen PM (1993) An annotated list of marine algae collected on Lanzarote Canary Islands January 1986. *Cour ForschungsInst Senckenberg* 159:93–102
- Kützing FT (1841) Über die "Polypieres calcifères" des Lamouroux. In: Kützing FT (ed) Zu der öffentlichen Prüfung sämmtlicher Classen der Realschule zu Nordhausen. 1841. Nordhausen, Realshule, pp 3–34
- Kützing FT (1849) Species algarum. Brockhaus, Lipsiae [Leipzig]
- Kützing FT (1858) Tabulae phycologicae oder Abbildungen der Tange, vol 8. Nordhausen
- Lamarck JB (1815) Suite des polypiers corticifère. *Mus Hist nat Paris* 2:76–84, 157–164, 227–240
- Lamouroux JV (1816) Histoire des polypiers coralligènes flexibles vulgairement nommés zoophytes. De l'imprimerie de F, Poisson, Caen
- Lamouroux JV (1821) Exposition méthodique des genres de l'ordre des Polypiers. Veuve Agasse, Paris
- Lemoine P (1924) Corallinacées du Maroc (1). *Bull Sci nat Maroc* 4:113–134
- Lemoine P (1929) Melobésières. *Biol Meddr K danske Vidensk Skr* 8:1–97
- Levrin T (1974) The marine algae of the archipelago of Madeira. *Bol Mus munic Funchal* 28:5–111
- Linnaeus C (1758) *Systema naturae per regna tria naturae*, vol 1. Stockholm
- López Hernández M, Gil-Rodríguez MC (1981) Estudio de la vegetación fisiológica del litoral comprendido entre Cabezo del Socorro y Montaña de la Mar, Güímar, Tenerife. *Vieraea* 11:141–170
- McNeill J, Barrie FR, Burdet HM, Demoulin V, Hawksworth DL, Marhold K, Nicolson DHJ, Prado J, Rado J, Silva PC, Skog JE, Wiersema JH, Turland NJ (2006) International Code of Botanical Nomenclature (Vienna Code) adopted by the 17th International Botanical Congress Vienna, Austria, July 2005. Gantner Verlag, Ruggell, Liechtenstein. (Note: Regnum Vegetabile, vol 146)
- Medina M, Haroun RJ (1993) Preliminary study on the dynamics of *Cystoseira abies-marina* populations in Tenerife (Canary Islands). *Cour ForschungsInst Senckenberg* 159:109–112
- Montagne JFC (1846) Ordo I. Phycaceae Fries. In: Durieu De Maisonneuve MC (ed) Durieu de Maisonneuve Flore d'Algérie. Cryptogamie. Première Partie. Imprimerie Royale, publiée par ordre du Gouvernement et avec le concours d'une Commission Académique, Paris, pp 1–197
- Montañés MA, Reyes J, Sansón M (2003) La comunidad de epífitos de *Zonaria tournefortii* en el norte de Tenerife (islas Canarias): análisis florístico y comentarios sobre su epifauna. *Vieraea* 31:121–132
- Moura CWN, Guimarães SMPB (2002) O gênero *Cheilosporum* (Decaisne) Zanardini (Corallinales, Rhodophyta) no litoral do Brasil. *Rev bras Biot* 25:65–77
- Muñoz E, Reyes J, Sansón M (2007) Descripción y cartografía de las comunidades bentónicas de Playa de San Marcos (Tenerife islas Canarias). *Vieraea* 35:93–108
- Neto AI (1994) Checklist of the benthic marine macroalgae of the Azores. *Arquipélago* 12A:15–34
- Neto AI (2000a) Ecology and dynamics of two intertidal algal communities on the littoral of the Island of São Miguel (Azores). *Hydrobiologia* 432:135–147
- Neto AI (2000b) Observations on the biology and ecology of selectes macroalgae from the littoral of São Miguel (Acores). *Bot Mar* 43:483–498
- Neto AI (2001) Macroalgal species diversity and biomass of subtidal communities of São Miguel (Azores). *Helgol Mar Research* 55:101–111
- Neto AI, Tittley I (1995) Structure and zonation of algal turf communities on the Azores: a numerical approach. *Bolm Mus munic Funchal Suppl* 4:487–504
- Neto AI, Cravo DC, Haroun RJ (2001) Checklist of the benthic marine plants of the Madeiran Archipelago. *Bot Mar* 44:391–414
- Newton L (1931) A handbook of the British seaweeds. British Museum (Natural History), London
- Norris JN, Johansen HW (1981) Articulated coralline algae of the Gulf of California México I: *Amphiroa* Lamouroux. *Smithson Contr Mar Sci* 9:i–iii + (1) + 1–29
- Otero-Schmitt J (1993) Some local patterns of zonation of benthic marine flora and fauna in Sal. Santiago, S. Vicente and Brava (Cape Verde Islands). *Cour ForschungsInst Senckenberg* 159:45–52
- Otero-Schmitt J, Sanjuan A (1992) Epibiotic seaweeds of the Cape Verde Islands. *Bot Mar* 35:379–390
- Palminha F (1957) Sobre a existencia de *Lithophyllum tortuosum* (Esper.) Foslie (= *Tenarea tortuosa* (Esper.) Lem.) nos Açores. *Bol Soc Port Ciênc Nat* 22:61–67
- Pérez-Cirera JL (1975) Catálogo florístico de las algas bentónicas de la Ría de Corme y Lage, NO. de España. *Anales Inst Bot Cavanilles* 32:5–87
- Piccone A (1884) Croceira del Corsaro alle isole Madera e Canarie del capitano Enrico d'Albertis. *Alge. Nuovo G bot Ital* 16:1–60
- Piccone A (1886) Alge del viaggio di circumnavigazione della Vettor Pisani. *Tipografia del Reale Istituto Sordo-Muti, Genova*
- Piccone A (1889) Alge della Crociera del alle Azzorre. *Nuovo G bot Ital Nuovo G bot Ital* 21:171–214
- Pinedo S, Sansón M, Afonso-Carrillo J (1992) Algas marinas bentónicas de Puerto de la Cruz (antes Puerto Orotava) Tenerife (Islas Canarias). *Vieraea* 21:29–60
- Porter DM (1987) Darwin's notes on Beagle plants. *Bull Br Mus nat Hist (Bot)* 14:145–233
- Price JH, John DM, Lawson GW (1986) Seaweeds of the western coast of tropical Africa and adjacents islands: a critical assessment. IV Rhodophyta (Florideae) 1. Genera A–F. *Bull Br Mus nat Hist (Bot)* 15:1–122
- Price JH, John DM, Lawson GW (1992) Seaweeds of the western coast of tropical Africa and adjacent islands: a critical assessment. IV. Rhodophyta (Florideae) 3. Genera H–K. *Bull Br Mus nat Hist (Bot)* 22:123–146
- Prud'homme van Reine WF (1988) Phytogeography of seaweeds of the Azores. *Helgoländer Meeresunters* 42:165–185
- Prud'homme van Reine WF, Haroun RJ, Audiffred PAJ (1994) A reinvestigation of Macaronesian seaweeds as studied by A. Piccone with remarks on those studied by A. Grunow. *Nova Hedwigia* 58:67–121
- Reyes J, Afonso-Carrillo J (1993) Morphology and anatomy of *Mesophyllum canariense* (Corallinaeae Rhodophyta) from the Canary Islands. *Cour ForschungsInst Senckenberg* 159:127–132
- Reyes J, Sansón M (1991) Adiciones a la flora marina de la isla de El Hierro (Islas Canarias). *Vieraea* 20:71–81
- Reyes J, Ocaña O, Sansón M, Brito A (2000) Descripción de comunidades bentónicas infralitorales en la Reserva Marina de La Graciosa e islotes del Norte de Lanzarote (Islas Canarias). *Vieraea* 28:137–154

- Rojas-González B, Afonso-Carrillo J (2000) Morfología y distribución de *Boergesenella fructiculosa* en las islas Canarias (Rhodophyta Rhodomelaceae). *Vieraea* 28:127–136
- Rojas-González B, Afonso-Carrillo J (2002a) Morfología y distribución de *Lophosiphonia cristata* y *L. reptabunda* en las islas Canarias (Rhodophyta Rhodomelaceae). *Vieraea* 30:31–44
- Rojas-González B, Afonso-Carrillo J (2002b) Notas corológicas sobre algas marinas de las islas Canarias: adiciones a la flora marina de El Hierro. *Vieraea* 30:99–107
- Rojas-González B, Afonso-Carrillo J (2004) Morfología y distribución de *Dipterosiphonia dendritica*, *D. rigens* y *D. reversa* en las islas Canarias (Rhodophyta Rhodomelaceae). *Vieraea* 32:135–150
- Rosas-Alquicira EF, Riosmena-Rodríguez R, Neto AI (2010) The lectotypification and status of *Amphiroa cryptarthrodia* (Lithophylloideae, Corallinales). *Phycol Res* (in press)
- Sangil C, Sansón M, Afonso-Carrillo J (2003a) Algas marinas de La Palma (islas Canarias): novedades florísticas y catálogo insular. *Vieraea* 31:83–119
- Sangil C, Afonso-Carrillo J, Sansón M (2003b) Zonación del fitobentos en el litoral sureste y adiciones a la flora marina de La Palma (Islas Canarias). *Rev Acad Canar Cienc* 15:310–319
- Sangil C, Afonso-Carrillo J, Sansón M (2004) Zonación del fitobentos en el litoral Noroeste de La Palma (Isla Canarias). *Rev Acad Canar Cienc* 16:75–90
- Sangil C, Sansón M, Afonso-Carrillo J (2005) Comunidades de algas esciáfilas en charcos eulitorales (La Palma, Islas Canarias). *Vieraea* 33:323–333
- Sansón M, Reyes J, Afonso-Carrillo J (1991) Contribution to the seaweed flora of the Canary Islands: new records of Florideophyceae. *Bot Mar* 34:527–536
- Santos Guerra A (1972) Contribución al estudio de la flora de la isla de La Gomera. *Vieraea* 1:86–102
- Santos Guerra A, Acuña Gonzalez A, Wildpret de la Torre W (1970) Contribución al estudio de la flora marina de la isla de La Palma. *Cuad Bot Canaria* 9:20–29
- Sauvageau C (1912) A propos des *Cystoseira* de Banyuls et Guéthary. *Bull Str Biol Arcachon* 14:133–556
- Schmidt OC (1929a) Die marine Vegetation der Azoren (Vorläufiger Bericht.). *Hedwigia* 68:327–346
- Schmidt OC (1929b) Beiträge zur Kenntnis der Meeresalgen der Azoren II. *Hedwigia* 69:165–172
- Schmidt OC (1931) Die marine Vegetation der Azoren in ihren Grundzügen dargestellt. *Bibliotheca Bot* 24:1–116
- Schneider CW, Searles RB (1991) Seaweeds of the southeastern United States. Cape Hatteras to Cape Canaveral. Duke University Press, Durham
- Seoane-Camba J (1965) Estudios sobre las algas bentónicas en la costa sur de la Península Ibérica (litoral de Cádiz). *Investigación pesq* 29:3–216
- Silva PC, Basson PW, Moe RL (1996) Catalogue of the benthic marine algae of the Indian Ocean. Univ Calif Publs in Bot 79:1–1259
- Sonder G (1845) Nova algarum genera et species, quas in itinere ad oras occidentales Novae Hollandiae, collegit L. Priess, Ph. Dr. Botanische Zeitung 3:49–57
- South GR, Tittley I (1986) A checklist and distributional index of the benthic marine algae of the North Atlantic Ocean. Huntsman Marine Laboratory and British Museum (Natural History), St. Andrews and London
- Stafleu FA, Cowan RS (1988) Taxonomic Literature, vol. VII: W-Z. Bohn Scheltema and Holkema, Utrecht, The Netherlands
- Stengenga H, Bolton JJ, Anderson RJ (1997) Seaweed of the south African west coast. *Contr Bolus Herbarium* 18:1–655
- Taylor WR (1960) Marine algae of the Eastern Tropical and Subtropical coasts of the Americas. University of Michigan Press, Ann Arbor
- Tittley I, Neto AI (2000) A provisional classification of algal characterised rocky shore biotopes in the Azores. *Hydrobiologia* 440:19–25
- Tittley I, Neto AI (2005) The marine algal (seaweed) flora of the Azores: additions and amendments. *Bot Mar* 48:248–255
- Tittley I, Neto AI (1994) Expedition Azores 1989. Benthic marine algae (seaweeds) record from Faial and Pico. *Arquipélago* 12A:1–13
- Tittley I, Neto AI, Farnham WF (1998) Marine algae of the island of Flores Azores: ecology and floristics. *Bolm Mus munic Funchal* 5:463–479
- Toste MF, Parente MI, Neto AI, Fletcher RL (2003) Life history of *Colpomenia sinuosa* (Scytoniphonaceae, Phaeophyceae) in the Azores. *J Phycol* 39:1268–1274
- Tuya F, Haroun RJ (2006) Spatial patterns and response to wave exposure of shallow water algal assemblages across the Canarian Archipelago: a multi-scaled approach. *Mar Ecol Prog Ser* 311:15–28
- Vickers A (1896) Contribution à la flore algologiques des Canaries. *Annls Sci nat (Bot)*, Ser 4:293–306
- Viera-Rodríguez MA (1987) Contribución al estudio de la flórula bentónica de la isla de La Graciosa Canarias. *Vieraea* 17:237–259
- Viera-Rodríguez MA, Wildpret de la Torre W (1986) Contribución al estudio de la vegetación bentónica de la isla de La Graciosa. *Vieraea* 16:211–231
- Viera-Rodríguez MA, Gil-Rodríguez MC, Audiffred PAJ, Prud'homme Van Reine WF, Haroun-Tabraue R, Wildpret de la Torre W (1987) Contribución al estudio de la flórula bentónica del islote de Montaña Clara. *Canarias*. *Vieraea* 17:271–279
- Villena-Balsa M, Afonso-Carrillo J, Wildpret de la Torre W (1987) Morfología, estructura y reproducción de una especie epífita del género *Jania* (Corallinaceae, Rhodophyta) de las Islas Canarias. *Vieraea* 17:19–42
- Walker RH, Brodie J, Russell S, Irvine LM, Orfanidis S (2009) Biodiversity of coralline algae in the Northeastern Atlantic including *Corallina caespitosa* sp. nov. (Corallinoideae, Rhodophyta). *J Phycol* 45:287–297
- Wallenstein FM, Neto AI (2006) Intertidal rocky shore biotopes of the Azores: a quantitative approach. *Helgol Mar Res* 60:196–206
- Weber-van Bosse A (1904) Corallinae verae of the Malay Archipelago II. In: Weber-van Bosse A, Foslie M (eds) The Corallinaceae of the Siboga-Expedition, vol 61. Siboga-Expeditie: ...Uitkomsten op Zoologisch, Botanisch, Oceanographisch en Geologisch Gebied... Monographie, pp 78–110
- Weisscher FCM (1982) Marine algae from Ilhéu de For a (Salvage Island) (Cancap-Project Contributions No. 13). *Bolm Mus munic Funchal* 34:23–34
- Weisscher FCM (1983) Marine algae from Selvagem Pequena (Salvage Island). *Bolm Mus munic Funchal* 35:41–80
- Woelkerling WJ, Nelson WA (2004) A baseline summary and analysis of the taxonomic biodiversity of Coralline red algae (Corallinales Rhodophyta) recorded from the New Zealand region. *Cryptogamie Algol* 25:39–106
- Woelkerling WJ, Sartoni G, Boddi S (2002) *Paulsivalvella huveorum* gen. et sp. nov. (Corallinaceae, Rhodophyta) from the Holocene of Somalia and Kenya, with a reassessment of *Lithothrix antiqua* from the Late Pleistocene of Mauritius. *Phycologia* 41:358–373
- Woelkerling WJ, Millar AJK, Harvey A, Baba M (2008) Recognition of *Pachyarthron* and *Bossiella* as distinct genera in the Corallinaceae, subfamily Corallinoideae (Coarlliinales, Rhodophyta). *Phycologia* 47:265–293
- Womersley HBS, Johansen HW (1996) Subfamily Corallinoideae. In: Womersley HBS (ed) The Marine benthic flora of Southern Australia—Part IIIB. Gracilariales Rhodymeniales Corallinales

- and Bonnemaisonales. Canberra Australian Biological Resources Study, Australia, pp 288–317
- Yendo K (1905) A revised list of Corallinae. J Coll Sci Imp Univ Tokyo 20:1–46
- Yoshida T (1998) Marine algae of Japan. Uchida Rokakuho Publishing, Tokyo
- Zanardini G (1840) (Lettera) alla direzione della biblioteca Italiana. Biblioteca Ital Milano 96:131–137
- Zanardini G (1844) Corallinee. Encycl. Ital. (Venezia:Tasso) 6:1013–1036