

Geographical variation in shell shape of the pod razor shell *Ensis siliqua* (Bivalvia: Pharidae)

Marta M. Rufino · Paulo Vasconcelos · Fábio Pereira ·
Juan Fernández-Tajes · Susana Darriba · Josefina Méndez ·
Miguel B. Gaspar

Received: 3 November 2011 / Revised: 14 March 2012 / Accepted: 27 March 2012 / Published online: 24 April 2012
© Springer-Verlag and AWI 2012

Abstract The present study assessed the existence of variation in the shell shape of the pod razor shell (*Ensis siliqua*) throughout its distributional range in the north-eastern Atlantic. Shells of *E. siliqua* caught at seven collecting sites (three in Portugal, three in Spain and one in Ireland) were studied by geometric morphometric methods, using both landmark- and contour-based methods. Both approaches (landmarks inside the valves and shell outline) discriminated the shells from Aveiro (centre of Portugal) and Strangford Lough (Ireland) from those caught in the nearby localities (remaining Portuguese and Spanish sites,

maximum distance of 550 km by sea). Landmark analysis revealed that shells from Aveiro were more similar to shells from Ireland (~1,500 km far away). Contour analysis revealed that shells from Aveiro had a shape with a comparatively larger height-to-width ratio, whereas shells from Ireland showed a slightly more curved outline than in the remaining sites. Landmark- and contour-based methods provided coherent complementary information, confirming the usefulness of geometric morphometric analyses for discerning differences in shell shape among populations of *E. siliqua*. A brief review of previous applications of geometric morphometric methods to modern bivalve species is also provided.

Communicated by Heinz-Dieter Franke.

M. M. Rufino (✉)
Centro Interdisciplinar de Investigação Marinha e Ambiental
(CIIMAR), Laboratório Nacional de Energia e Geologia
(LNEG), Departamento de Geologia Marinha (DGM),
Estrada da Portela, Zambujal, Apartado 7586,
2721-866 Alfragide, Portugal
e-mail: marta.m.rufino@gmail.com

P. Vasconcelos · F. Pereira · M. B. Gaspar
Instituto Nacional de Recursos Biológicos (INRB, I.P.)/IPIMAR,
Avenida 5 de Outubro s/n, 8700-305 Olhão, Portugal

P. Vasconcelos
Centro de Estudos do Ambiente e do Mar (CESAM),
Departamento de Biologia, Universidade de Aveiro,
Campus de Santiago, 3810-193 Aveiro, Portugal

J. Fernández-Tajes · J. Méndez
Departamento de Biología Celular y Molecular,
Universidad da A Coruña, A Zapateira s/n,
15071 Coruña, Spain

S. Darriba
Instituto Tecnológico para o Control do Medio Mariño de
Galicia (INTECMAR), Peirao de Vilaxoán s/n,
36611 Vilagarcía de Arousa, Pontevedra, Spain

Keywords Pod razor shell · *Ensis siliqua* · Shell shape ·
Geometric morphometrics

Introduction

The comparison of anatomical features of organisms has been a central element in biology for centuries (Adams et al. 2004). Shell morphology, which preserves the ontogenetic record of growth, is now the principal subject of a subfield of morphometrics (Madec et al. 2003), namely theoretical morphology, which since the initial mathematical modelling of Raup (1966) has led to convincing studies on the formal and historical determinants of shell forms, as well as functional interpretations of their observed distributions in theoretical morphospaces (e.g. Stone 1996, 1999; McGhee 1999; Samadi et al. 2000). Several models of theoretical morphology of molluscan shells have been successfully developed in previous studies (Savazzi et al. 1982; Savazzi 1989, 1990; McClain et al. 2004). In more traditional approaches, the shape of molluscan shells has

also been characterised by using simpler measurements, such as the ratio of width to height (for a review see Goodfriend 1986), or by plotting these two measurements in order to create a “size/shape space” (Cain 1981; Cameron and Cook 1989). More recently, geometric morphometric (GM) methods allowed the partitioning of shape and size components, preserving the main geometric properties of the specimens and generating a visual representation and shape variables than can be statistically analysed. GM is often described as a “revolution” in the area of morphometrics (Adams et al. 2004) and has been successfully applied in most fields of science, since these techniques are useful for biology, ecology, evolution, ontogeny, taxonomy, among other studies.

Due to their hard un-deformable shells, bivalves are an excellent group to apply GM methods (Rufino et al. 2007). Historically, GM techniques have been developed within two main approaches: (1) the analysis of outline or contour, where the frontier/border of a specimen is analysed (Rohlf and Archie 1984) and (2) the analysis of landmarks, where the change in position of homologous points identified in the specimens is quantified. Recent studies use a mixture of these two techniques, called semi-landmarks. Both approaches provide different types of information and can be applied in distinct circumstances. The first GM studies with bivalves used mainly contour-based methods (Ferson et al. 1985; Innes and Bates 1999), whereas presently most works employ landmarks or a combination of both approaches, which have been considered to be more powerful (Rufino et al. 2007). These techniques have been successfully applied to distinguish similar species of bivalves (Rufino et al. 2007; Costa et al. 2010), between fossil and modern specimens (Aguirre et al. 2006; Gordillo et al. 2011) or between wild and aquaculture stocks (Valladares et al. 2010), to detect ontogenetic shape changes (Márquez et al. 2010a) and to analyse geographical variation in shape (Palmer et al. 2004; Sousa et al. 2007; Costa et al. 2008; Márquez et al. 2010b; Gordillo et al. 2011; Márquez and Van der Molen 2011). A compilation of previous works that applied GM methods to modern bivalves is summarised in Table 1.

The pod razor shell *Ensis siliqua* (Linnaeus, 1758) is distributed in the Mediterranean Sea and in the Atlantic Ocean, from Morocco to Norway (Tebble 1966), being commercially exploited in Portugal, Spain, Ireland and UK (Guerra and Lodeiros 2008). The most common methods used to harvest *E. siliqua* are hand-gathering (Fahy and Carroll 2007; Constantino et al. 2009), and mechanic or hydraulic dredging (Gaspar and Monteiro 1998; Robinson and Richardson 1998; Gaspar et al. 1999; Tuck et al. 2000; Fahy and Gaffney 2001; Hauton et al. 2003a, b). This infaunal species inhabits the low intertidal zone down to

40 m depth (Fahy and Gaffney 2001; Rufino et al. 2010), occurring in substrates of fine to medium sand. It lives burrowed into the sediment with the long axis of the shell orientated roughly vertical, extending the siphons out of the surface to feed on suspended particles (Gaspar and Monteiro 1998; Gaspar et al. 1999; Fahy and Gaffney 2001; Rufino et al. 2010). The pod razor shell has a large and powerful foot that enables vertical burrowing into sand up to 60 cm depth. The characteristic shape of the shell of *E. siliqua*, elongated, laterally compressed, narrowed and with almost semi-cylindrical valves, facilitates rapid burrowing.

Most previous studies with *E. siliqua* have focused on life cycle and growth (Henderson and Richardson 1994; Gaspar and Monteiro 1998; Fahy and Gaffney 2001; Darriba et al. 2005; da Costa et al. 2010), larval cycle and aquaculture potential (Darriba et al. 2005; da Costa et al. 2010), genetics (Fernández-Tajes et al. 2007; Varela et al. 2007; Freire et al. 2008; Arias et al. 2011) or on fishery-related aspects (Robinson and Richardson 1998; Gaspar et al. 1999; Tuck et al. 2000; Fahy and Gaffney 2001; Hauton et al. 2003a, b; Constantino et al. 2009). To the author’s best knowledge, the shell shape of *E. siliqua* has never been analysed by applying GM approaches. Therefore, the present study aimed to analyse and compare the shape of *E. siliqua* shells across its distributional range in the north-eastern Atlantic (Portugal, Spain and Ireland), using both landmark- and contour-based methods, in order to assess the suitability of these techniques for discerning populations of this species.

Materials and methods

Study area and sample collection

Individuals of *E. siliqua* with a shell length ranging from 12 to 15 cm were caught by scuba diving or by dredging at seven collecting sites: three sites in Portugal (Olhão—PO, Setúbal—PS and Aveiro—PA), three sites in Spain (Barra de Cangas—SB, Fisterra—SF and Celeiro—SC) and one site in Ireland (Strangford Lough—IS) (Fig. 1). The southernmost and northernmost collecting sites are separated by approximately 17 degrees of latitude (Olhão $\approx 37^\circ\text{N}$ and Strangford Lough $\approx 54^\circ\text{N}$), corresponding to a distance of almost 2,000 km (more than 1,000 nmi). In the laboratory, the soft parts were removed and the valves were cleaned, air-dried and stored for subsequent analyses.

In order to decrease unpredictable sources of variability, only the right valve of each shell was used for GM purposes. A preliminary analysis of *E. siliqua* shells revealed that the muscle scars on the inner surface of the valves

Table 1 Compilation of previous studies that applied geometric morphometric methods to modern bivalves

Species	Country or geographical area	Type/subject	Contour/landmarks	Reference
<i>Mytilus</i> sp.			Elliptic Fourier analysis	Ferson et al. (1985)
<i>Mytilus edulis</i>	Canada (eastern Newfoundland) (16)	Multispecies	300–600 points, 10 harmonics	Innes and Bates (1999)
<i>Mytilus trossulus</i>				
<i>Chamelea gallina</i>	Spain (6), Italy (1)	Geographical	100 points, 10 harmonics	Palmer et al. (2004)
<i>Brachidontes darwinianus</i>	Argentina: Quaternary fossils (19), modern (8)	Fossil/modern multispecies geographical	1 landmark (umbo) + 39 semi-landmarks	Aguirre et al. (2006)
<i>Brachidontes purpuratus</i>				
<i>Brachidontes rodriguezii</i>				
<i>Mytilus chilensis</i>	Chile (8)	Geographical	100 points, 10 harmonics	Krapivka et al. (2007)
<i>Chamelea gallina</i>	Portugal (1)	Multispecies	10 harmonics + 9 landmarks	Rufino et al. (2007)
<i>Chamelea striatula</i>				
<i>Corbicula fluminea</i>	Portugal (6)	Geographical	11 landmarks	Sousa et al. (2007)
<i>Ruditapes decussatus</i>	Italy (9), Spain (1), Tunisia (1)	Multispecies geographical	7 harmonics	Costa et al. (2008)
<i>Ruditapes philippinarum</i>				
<i>Anomalocardia brasiliana</i>	Thailand (7), Philippines (1), Jamaica (1), Colombia (1), Brazil (1), Ecuador (1)	Multispecies geographical	19 landmarks	Roopnarine et al. (2008)
<i>Anomalocardia squamosa</i>				
<i>Ilioichione subrugosa</i>				
<i>Mytilus edulis</i>	Canada (7), USA (1), UK (5), France (1), Finland (1)	Multispecies geographical	300–600 points, 10 harmonics	Gardner and Thompson (2009)
<i>Mytilus galloprovincialis</i>				
<i>Mytilus trossulus</i>				
<i>Ruditapes decussatus</i>	Italy (2)	Multispecies	180 points, 7 harmonics	Costa et al. (2010)
<i>Ruditapes philippinarum</i>				
<i>Aequipecten tehuelchus</i>	Argentina (1)	Ontogenetic	8 landmarks + 20 semi-landmarks	Márquez et al. (2010a)
<i>Ameghinomya antiqua</i>	Argentina (5)	Geographical	10 harmonics + 13 landmarks	Márquez et al. (2010b)
<i>Mytilus chilensis</i>	Chile (6)	Geographical wild versus cultivated	9 landmarks	Valladares et al. (2010)
<i>Tawera gayi</i>	Chile (3), Argentina (2), New Zealand (1), South Africa (1)	Fossil/modern multispecies geographical	10 harmonics	Gordillo et al. (2011)
<i>Tawera philomela</i>				
<i>Tawera spissa</i>				
<i>Ensis macha</i>	Argentina (12)	Geographical	15 harmonics	Márquez and Van der Molen (2011)
<i>Pectinidae</i>	USA	Multispecies	Sliding semi-landmarks in 3D	Serb et al. (2011)

The number of sites or populations analysed is indicated between brackets. In contour and landmark analyses, the number of points or harmonics and the number of landmarks are also indicated

were very difficult to identify, and thus to increase precision, shells were examined under a binocular microscope and the selected landmarks were carefully marked with a pen. Each valve was then digitised using a previously calibrated HP® Scanjet 5530 before subsequent GM analyses. The number of valves subjected to landmark and contour analyses differed due to the adequacy of each valve for the method employed. Indeed, valves with slightly damaged margins can be used for landmark analysis but not for contour analysis.

Landmark analysis

A total of 334 individuals were used for landmark analysis: 159 valves from Portugal (PO = 112, PS = 28 and PA = 19), 119 valves from Spain (SB = 49, SF = 60 and SC = 10) and 56 valves from Ireland (IS = 56). In each valve, 11 shell landmarks were selected to quantify shape: the first, second and third landmarks were located at the pallial sinus, the fourth and fifth in the limits of the scar of the posterior adductor muscle, the sixth and seventh

Fig. 1 Geographical location of the collecting sites of the pod razor shell (*Ensis siliqua*) in Portugal (Olhão—*PO*, Setúbal—*PS* and Aveiro—*PA*), Spain (Barra de Cangas—*SB*, Fisterra—*SF* and Celeiro—*SC*) and Ireland (Strangford Lough—*IS*)




Fig. 2 Location and description of the landmarks in the inner surface of the shell of *Ensis siliqua*: (1) ventralmost point of the pallial sinus, (2) centralmost point (maximum curvature) of the pallial sinus, (3) dorsalmost point of the pallial sinus, (4) posteriormost point of the posterior adductor muscle, (5) anteriormost point of the posterior adductor muscle, (6) dorsalmost point of the posterior limit of the

anterior adductor muscle, (7) ventralmost point of the posterior limit of the anterior adductor muscle, (8) ventral anterior limit of the pallial line, (9) dorsal anterior limit of the pallial line, (10) dorsalmost point of the anterior margin of the valve (pseudo-landmark), (11) ventralmost point of the anterior margin of the valve (pseudo-landmark)

corresponded to the posterior limits of the scar of the anterior adductor muscle, the eighth and ninth were located at the anterior limits of pallial line, whereas the tenth and eleventh (pseudo-landmarks) were located in the anterior margin of the valve (Fig. 2). Because most valves analysed were damaged in the posterior margin, no landmarks were selected in this area of the shell.

A more detailed description of GM methods using landmarks has been reported by Adams et al. (2004); therefore, only a brief explanation is provided here. After digitisation, landmark positions were rotated, scaled (to

unit centroid size) and translated through Generalized Procrustes Analysis (GPA), a procedure that eliminates variation due to differences in scale, size and orientation. An average map (consensus configuration) of the shell shape, plus the uniform and non-uniform (partial warps) components of shape variation were calculated. The two uniform components describe differences that affect equally all parts of the shell (global differences). The non-uniform components (partial warps) describe localised departures from the average map of the shell. Partial warps together with uniform component are the statistical

variables of shell shape. The effect of size (allometry) was tested through permutation bifactorial multivariate ANOVA applied to partial warps and uniform component, treating centroid size and geographical location (collecting site) as independent variables.

To check whether the shape variables were normally distributed, Shapiro–Wilk test for multivariate normality was performed (mvnormtest package; Jarek 2009), followed by permutational multivariate analysis of variance (adonis, vegan package; Oksanen et al. 2010), to determine whether there were significant differences depending on geographical location (collecting site) or whether there was a significant allometric effect (i.e. shape changes with size, quantified as centroid size). Relative warp analysis, a technique analogous to principal component analysis, was applied to determine the patterns of shape variation. The mean landmark configuration of the extremes of the two principal relative warps was estimated to determine which landmarks were varying. Coordinates of the landmarks were calculated from the digitised images of the shells using tpsDig (Rohlf 2003a). Partial warps and centroid size were calculated using tpsReg (Rohlf 2000), and relative warps were estimated using tpsRel (Rohlf 2003b). All other statistical analyses and graphical representations were performed using the free software R (R Development Core Team 2010).

Contour analysis

A total of 224 individuals were examined for contour analysis: 128 valves from Portugal (PO = 86, PS = 27 and PA = 15), 66 valves from Spain (SB = 26, SF = 33 and SC = 7) and 30 valves from Ireland (IS = 30). The overall shell shape was studied through elliptic Fourier analysis of the contour coordinates (Rohlf and Archie 1984). For each shell, 100 coordinates of the outline were obtained using tpsDig (Rohlf 2003a). In a preliminary study based on visual inspection, it was concluded that 15 harmonics were sufficient to properly characterise the shell shape of *E. siliqua*, similar to the study by Márquez and Van der Molen (2011) with *E. macha*. As recommended by Rohlf and Archie (1984), 59 elliptic Fourier coefficients were normalised mathematically to be invariant of size, location, rotation and starting position (which was always approximately the left lower extreme of the shell). All further Fourier analyses were performed using the software Morpheus (Slice 1998). The mean outline of the shells of *E. siliqua* from each collecting site was reconstructed using Fourier analysis. The Shapiro–Wilk test for multivariate normality was applied to the harmonic coefficients to verify whether the data presented a Gaussian multivariate distribution (mvnormtest package; Jarek 2009), followed by permutational multivariate

analysis of variance (adonis, vegan package; Oksanen et al. 2010).

Results

Landmark analysis

After a detailed preliminary analysis, 11 landmarks were found essential to properly describe the main shape features of the shells of *E. siliqua*. Centroid size of the individuals differed between collecting sites (Fig. 3), with a north–south decreasing trend in mean shell length, which was largest in *E. siliqua* from Strangford Lough (IS) and smallest in Olhão (PO) (Kruskal–Wallis rank sum test: $H = 179.727$, $P < 0.001$). Still, there was some overlap in sizes across all collecting sites, which allowed both to compare the shape of the bivalves and to test for an eventual allometric effect.

The shape variables were not normally distributed (Shapiro–Wilks test for multivariate normality: $W = 0.764$, $P < 0.001$) and the variances differed significantly between collecting sites (Anderson’s multivariate homogeneity of group dispersions: ANOVA: $F = 18.208$, $P < 0.001$; Anderson 2006), with the shells from Olhão (PO) showing greater dispersions than the remaining samples. Therefore, nonparametric MANOVA was used to test for shape differences (Anderson 2001) between collecting sites, using centroid size as covariable. The nonparametric MANOVA indicated that both the covariable




Fig. 3 Centroid size (square root of the sum of squared distances of a set of landmarks from their centroid as an accurate measure of size) in the shells of *Ensis siliqua* from each collecting site. The central dot represents the median, the limits of the box, the first and third quartiles, and the whiskers, the 95 % confidence interval. Abbreviations: Olhão—PO, Setúbal—PS, Aveiro—PA, Barra de Cangas—SB, Fisterra—SF, Celeiro—SC and Strangford Lough—IS

centroid size ($r^2 = 0.011$, $F = 5.182$, $P = 0.003$) and the collecting site ($r^2 = 0.254$, $F = 18.832$, $P = 0.001$) were significantly different, whereas their interaction was not statistically significant ($r^2 = 0.015$, $F = 1.136$, $P = 0.286$). In practice, this means that there is a significant change in shape with size (allometry), as well as significant differences in shape between collecting sites, but the change in shape with specimen size is similar for all studied populations.

Relative warp analysis showed clear trends in shape change, although with overlap of points in all collecting sites, typical in this kind of morphometric data (Fig. 4). The first three relative warps (RW1, RW2 and RW3) explained 50.4, 13.8 and 5.8 % of shape variability, respectively. The RW1 shows the main axis of shape change, distinguishing the shells from Aveiro (PA) and Strangford Lough (IS) from those of Olhão (PO), with the remaining populations being intermediate. The reconstruction of shape variation on this axis allowed detecting that the main changes occurred in the distance between landmark 3 and both landmarks 4–5 and 6–7, which was smaller in PA and IS than in shells from other collecting sites. In practice, this means that *E. siliqua* from PA and IS

have the pallial sinus (landmark 3) closer to the scars of the posterior (landmarks 4 and 5) and anterior adductor muscles (landmarks 6 and 7) than *E. siliqua* from PO. Furthermore, these shells from Olhão (PO) presented two peaks of shape distribution over the RW1, which had no correspondence with centroid size, indicative of some morphological plasticity in this population. The RW2 detected some differences between the shells from Aveiro (PA, in the positive extreme) and the shells from Strangford Lough (IS, in the negative extreme), with the remaining populations being intermediate (Fig. 4). The shape reconstruction for the RW2 extremes revealed that shells in the positive side (where shells from PA are located) have greater distance between landmarks 4–5 and 6–7 than those in the negative side (where shells from IS are located). In practice, this means that *E. siliqua* from PA have a relatively longer body between the scars of the posterior (landmarks 4 and 5) and anterior adductor muscles (landmarks 6 and 7) than those from IS, with the remaining populations being intermediate between these two morphs.

The linear model of the RW1 with centroid size and collecting site showed that the main effects were both




Fig. 4 Relative warp analysis of the landmarks in the shells of *Ensis siliqua* from each collecting site and respective shape TPS reconstruction of main axis of variation. Abbreviations: Olhão—PO,

Setúbal—PS, Aveiro—PA, Barra de Cangas—SB, Fisterra—SF, Celeiro—SC and Strangford Lough—IS



Fig. 5 Mean contour of the shells of *Ensis siliqua* from each collecting site, reconstructed using Fourier analysis. Abbreviations: Olhão—PO, Setúbal—PS, Aveiro—PA, Barra de Cangas—SB, Fisterra—SF, Celeiro—SC and Strangford Lough—IS

significant (ANCOVA: F centroid = 8.881, $P = 0.003$; F site = 11.265, $P < 0.001$), although without significant interaction between them (ANCOVA: F interaction = 0.796, $P = 0.574$). Altogether, this revealed that the relationship between the RW1 and centroid size was not significantly different among collecting sites, thus not relevant in terms of shape change across this axis. Furthermore, the plot of the residuals (data omitted for brevity) showed similar results and confirmed this same trend.

Contour analysis

The harmonic coefficients were not normally distributed (Shapiro–Wilks test for multivariate normality: $W = 0.193$, $P < 0.001$) and the variances differed significantly between collecting sites (Anderson’s multivariate homogeneity of group dispersions: ANOVA $F_{[6, 217]} = 3.642$, $P = 0.001$; Anderson 2006), with the shells of *E. siliqua* from Setúbal (PS) and Strangford Lough (IS) showing greater dispersions than the remaining samples. The non-parametric MANOVA confirmed the existence of significant differences in the contour of shells between collecting sites ($r^2 = 0.227$, $F = 10.624$, $P < 0.001$). The mean shell outline in each collecting site in relation to the global mean revealed that *E. siliqua* from Aveiro (PA) have thicker and shorter shells than in the remaining collecting sites, whereas shells from Strangford Lough (IS) are slightly more curved (Fig. 5).

Discussion

Geometric morphometric methods applied in the study of *E. siliqua* shell morphology were able to discern three main shapes, corresponding to different geographical groups among seven collecting sites distributed along the north-eastern Atlantic coasts (Portugal, Spain and Ireland). As in previous studies, both landmark- and contour-based methods were used, because these provide different and complementary information, namely on the position of internal features of the shell (landmark analysis) that are related with physiological aspects (e.g. muscle scars), as well as on the shell outline (contour analysis). Still, both approaches

provided coherent complementary results, separating the populations of *E. siliqua* of Aveiro (Portugal) and Strangford Lough (Ireland) from the remaining collecting sites. Indeed, the first relative warp showed that the shells from Aveiro (PA) were more similar to those from Strangford Lough (IS), which is located more than 1,500 km apart (around 840 nmi), than to the shells from the other Portuguese (PO and PS) and Spanish (SB, SF and SC) sites, which are situated within a maximum distance slightly over 400 km (almost 220 nmi). The shells from Olhão (PO) extended throughout the first relative warp, denoting a certain degree of morphological plasticity in this southernmost population of *E. siliqua*. On the whole, combining the results gathered using both methods (landmark and contour analyses), three main groups with different shell shape were identified: one morph from Strangford Lough (IS), another from Aveiro (PA) and another comprising the remaining populations of *E. siliqua* from the Iberian Peninsula (Portugal: PO and PS; Spain: SB, SF and SC).

The results from landmark analysis are corroborated by recent genetic studies with *E. siliqua*, where the larger differences were observed between Iberian and Irish populations, except for Aveiro individuals that were closer to northern individuals (Arias et al. 2011). These authors argued that the Aveiro population might be predominantly self-recruiting, as a consequence from some mechanism that could affect larval dispersal of *E. siliqua*. A recent geophysical model, where particles simulating larvae were released into the main Portuguese estuaries, estimated an along shore dispersal distance around 60 km for crab zoeae and greater than 100 km for crab megalopae (Peliz et al. 2007). Furthermore, these authors concluded that larvae released in the Ria de Aveiro can easily reach northern estuaries up to Minho River, and only if these estuaries are also sources of larvae, these could reach Fisterra, but not northern than this location. These findings support the similarities in shell shape among the Spanish populations of *E. siliqua* (SB, SF and SC), but do not explain the differences in shape between the shells from Aveiro (PA) and from the other collecting sites along the Portuguese (PO and PS) and Spanish (SB, SF and SC) coasts. Peliz et al. (2007) also found little transport southern of the Mondego River, which excludes larval transport by along shore

currents as a factor responsible for the similarities observed between the shells of *E. siliqua* from Portuguese (PO and PS) and Spanish (SB, SF and SC) populations. Moreover, crab larvae have a longer pelagic larval duration than *E. siliqua*, which under laboratory conditions lasts for only 15–20 days before settlement (Darriba et al. 2005; da Costa et al. 2010). Probably for this reason, the pod razor shell is a typical inhabitant of sandy beaches, whose larvae find settlement spots in the vicinities of the source populations along the coasts of the Iberian Peninsula.

As mentioned above, there was a recent study focusing on the genetic identification of *E. siliqua* populations along the north-eastern Atlantic (Arias et al. 2011). However, the phenotype of an organism does not always directly reflect its genotype information, and often genetic evidence does not match with results from shape analysis (Lawing and Polly 2009). In addition, results from genetic studies greatly depend on the methods applied and on the markers used. For instance, Sousa et al. (2007) studying an invasive bivalve in two Portuguese estuaries were able to identify morphological differences through GM methods, where no genetic evidences were observed. Indeed, GM is an excellent tool to identify phenotypic differences related to environmental specificities in different locations and thus ideal to study geographical shape changes. Arias et al. (2011) considered that genetic similarities between populations of *E. siliqua* from Aveiro (Portugal) and Ireland are most likely due to anthropogenic influence, for instance through larval transport in ballast water of ships. However, although the existence of an important commercial harbour, a shipyard and a fishing port in the Ria de Aveiro, routes connecting Aveiro and Ireland are certainly sporadic.

Morphometrics is an evolving discipline, providing increasingly powerful techniques for quantitative characterisation and comparison of shape (Roth and Mercer 2000). Before the 1990s, morphometric tools were restricted to bivariate plots and multivariate statistics (Urdu et al. 2010a), but since then enhancements included the development of GM that more directly preserve geometry and the spatial relationships among landmark points and the contours of a form (Roth and Mercer 2000). Bivalves have hard shells, which make them excellent candidates for geographical studies of morphological variation in individuals from different populations, based solely on the analysis of their shape profiles (Costa et al. 2008). In the present study, the use of two complementary GM methods (landmark and contour analysis) was fundamental. Shape differences detected in *E. siliqua* caught at several collecting sites were not limited to the shell outline, but also occurred in the inner surface of the valves, that is reflecting differences in the internal structure of the bivalves. Variation in mollusc shell morphology can be both genetically and environmentally determined. However, although there

is an extensive literature on correlations between some environmental aspects and mollusc shell shapes (Urdu et al. 2010b), the contribution of local adaptation to morphological differentiation of bivalve populations is still poorly studied (Costa et al. 2008). In this context, further research should focus on the phenotypic variation and morphological plasticity of the shell of *E. siliqua* as a function of different environmental conditions (e.g. hydrodynamics, depth, sediment type and grain size). Altogether, these environmental variables are known to influence bivalve's burrowing ability and thus might also play a key role in the geographical variation in shell shape of *E. siliqua*.

Acknowledgments The authors would like to thank Dr. Dai Roberts and Adele Cromie for providing samples of pod razor shells from Ireland. This study was funded by Community Initiative Programmes (INTERREG-IIIb, Atlantic Area) Sustainable HARvesting of Ensis (090-SHARE) and Towards Integrated Management of Ensis Stocks (206-TIMES) from the European Community. Marta M. Rufino and Paulo Vasconcelos benefited from postdoctoral grants (SFRH/BPD/14935/2004 and SFRH/BPD/26348/2006, respectively) awarded by the Fundação para a Ciência e Tecnologia (FCT—Portugal). Finally, the authors acknowledge three anonymous referees for valuable comments and suggestions that greatly improved the revised manuscript.

References

- Adams DC, Rohlf FJ, Slice D (2004) Geometric morphometrics: ten years of progress following the “revolution”. *Ital J Zool* 71:5–16
- Aguirre ML, Perez SI, Sirch YN (2006) Morphological variability of *Brachidontes Swainson* (Bivalvia, Mytilidae) in the marine Quaternary of Argentina (SW Atlantic). *Palaeogeogr Palaeoclimatol Palaeoecol* 239:100–125
- Anderson MJ (2001) A new method for non-parametric multivariate analysis of variance. *Austral Ecol* 26:32–46
- Anderson MJ (2006) Distance-based tests for homogeneity of multivariate dispersions. *Biometrics* 62:245–253
- Arias A, Fernández-Moreno M, Fernández-Tajes J, Gaspar MB, Méndez J (2011) Strong genetic differentiation among east Atlantic populations of the sword razor shell (*Ensis siliqua*) assessed with mtDNA and RAPD markers. *Helgol Mar Res* 65:81–89
- Cain AJ (1981) Variation in shell shape and size of helicid snails in relation to other pulmonates in faunas of the Palaearctic region. *Malacologia* 21:149–176
- Cameron RAD, Cook LM (1989) Shell size and shape in Madeiran land snails—do niches remain unfilled. *Biol J Linn Soc* 36:79–96
- Constantino R, Gaspar MB, Pereira F, Carvalho S, Cúrdia J, Matias D, Monteiro CC (2009) Environmental impact of razor clam harvesting using salt in Ria Formosa lagoon (southern Portugal) and subsequent recovery of associated benthic communities. *Aquat Conserv Mar Freshwat Ecosyst* 19:542–553
- Costa C, Aguzzi J, Menesatti F, Antonucci F, Rimatori V, Mattoccia M (2008) Shape analysis of different populations of clams in relation to their geographical structure. *J Zool* 276:71–80
- Costa C, Menesatti P, Aguzzi J, D'Andrea S, Antonucci F, Rimatori V, Pallottino F, Mattoccia M (2010) External shape differences between sympatric populations of commercial clams *Tapes decussatus* and *T. philippinarum*. *Food Bioprocess Technol* 3:43–48

- da Costa F, Martínez-Patiño D, Ojea J, Nóvoa S (2010) Larval rearing and spat production of the razor clam *Ensis siliqua* (Bivalvia: Pharidae). *J Shellfish Res* 29:347–351
- Darriba S, San Juan F, Guerra A (2005) Gametogenic cycle of *Ensis siliqua* (Linnaeus, 1758) in the Ría de Corcubión, northwestern Spain. *J Mollus Stud* 71:47–51
- Fahy E, Carroll J (2007) Consequences of hydraulic dredging for a razor clam *Ensis siliqua* (L.) bed in the north-west Irish Sea. *Biol Environ* 107B:115–128
- Fahy E, Gaffney J (2001) Growth statistics of an exploited razor clam (*Ensis siliqua*) bed at Gormanstown, Co Meath, Ireland. *Hydrobiologia* 465:139–151
- Fernández-Tajes J, Gaspar MB, Martínez-Patiño D, McDonough N, Roberts D, González-Tizón A, Martínez-Lage A, Méndez J (2007) Genetic variation of the razor clam *Ensis siliqua* (Jeffreys, 1875) along the European coast based on random amplified polymorphic DNA markers. *Aquacult Res* 38:1205–1212
- Ferson SF, Rohlf FJ, Koehn RK (1985) Measuring shape variation of two-dimensional outlines. *Syst Zool* 34:59–68
- Freire R, Fernández-Tajes J, Méndez J (2008) Identification of razor clams *Ensis arcuatus* and *Ensis siliqua* by PCR–RFLP analysis of ITS1 region. *Fish Sci* 74:511–515
- Gardner JPA, Thompson RJ (2009) Influence of genotype and geography on shell shape and morphometric trait variation among North Atlantic blue mussel (*Mytilus* spp.) populations. *Biol J Linn Soc* 96:875–897
- Gaspar MB, Monteiro CC (1998) Reproductive cycles of the razor clam *Ensis siliqua* and the clam *Venus striatula* off Vilamoura, southern Portugal. *J Mar Biol Assoc UK* 78:1247–1258
- Gaspar MB, Castro M, Monteiro CC (1999) Effect of tooth spacing and mesh size on the catch of the Portuguese clam and razor clam dredge. *ICES J Mar Sci* 56:103–110
- Goodfriend GA (1986) Variation in land-snail shell form and size and its causes—a review. *Syst Zool* 35:204–223
- Gordillo S, Márquez F, Cárdenas J, Zubimendi MÁ (2011) Shell variability in *Tavera gayi* (Veneridae) from southern South America: a morphometric approach based on contour analysis. *J Mar Biol Assoc UK* 91:815–822
- Guerra A, Lodeiros C (2008) Navajas y longueirones: biología, pesquerías y cultivo. Consellería de Pesca e Asuntos Marítimos, Xunta de Galicia, Santiago de Compostela
- Hauton C, Atkinson RJA, Moore PG (2003a) The impact of hydraulic blade dredging on a benthic megafaunal community in the Clyde Sea area, Scotland. *J Sea Res* 50:45–56
- Hauton C, Hall-Spencer JM, Moore PG (2003b) An experimental study of the ecological impacts of hydraulic bivalve dredging on maerl. *ICES J Mar Sci* 60:381–392
- Henderson SM, Richardson CA (1994) A comparison of the age, growth rate and burrowing behaviour of the razor clams, *Ensis siliqua* and *E. ensis*. *J Mar Biol Assoc UK* 74:939–954
- Innes DJ, Bates JA (1999) Morphological variation of *Mytilus edulis* and *Mytilus trossulus* in eastern Newfoundland. *Mar Biol* 133:691–699
- Jarek S (2009) mvnrmtest: normality test for multivariate variables. R package version 0.1-7
- Krapivka S, Toro JE, Alcapán AC, Astorga M, Presa P, Pérez M, Guiñez R (2007) Shell-shape variation along the latitudinal range of the Chilean blue mussel *Mytilus chilensis* (Hupe 1854). *Aquacult Res* 38:1770–1777
- Lawing AM, Polly PD (2009) Geometric morphometrics: recent applications to the study of evolution and development. *J Zool* 280:1–7
- Madec L, Bellido A, Guiller A (2003) Shell shape of the land snail *Cornu aspersum* in North Africa: unexpected evidence of a phylogeographical splitting. *Heredity* 91:224–231
- Márquez F, Van der Molen S (2011) Intraspecific shell-shape variation in the razor clam *Ensis macha* along the Patagonian coast. *J Mollus Stud* 77:123–128
- Márquez F, Amoroso R, Sainz MFG, Van der Molen S (2010a) Shell morphology changes in the scallop *Aequipecten tehuelchus* during its life span: a geometric morphometric approach. *Aquat Biol* 11:149–155
- Márquez F, Robledo J, Peñaloza GE, Van der Molen S (2010b) Use of different geometric morphometrics tools for the discrimination of phenotypic stocks of the striped clam *Ameghinomya antiqua* (Veneridae) in north Patagonia, Argentina. *Fish Res* 101:127–131
- McClain CR, Johnson NA, Rex MA (2004) Morphological disparity as a biodiversity metric in lower bathyal and abyssal gastropod assemblages. *Evolution* 58:338–348
- McGhee GR Jr (1999) Theoretical morphology. Columbia University Press, New York
- Oksanen J, Blanchet FG, Kindt R, Legendre P, O’Hara RB, Simpson GL, Solymos P, Stevens MHH, Wagner H (2010) vegan: community ecology package. R package version 1.17-4. <http://CRAN.R-project.org/package=vegan>
- Palmer M, Pons GX, Linde M (2004) Discriminating between geographical groups of a Mediterranean commercial clam (*Chamelea gallina* (L.): Veneridae) by shape analysis. *Fish Res* 67:93–98
- Peliz A, Marchesiello P, Dubert J, Marta-Almeida M, Roy C, Queiroga H (2007) A study of crab larvae dispersal on the Western Iberian Shelf: physical processes. *J Mar Syst* 68:215–236
- Raup DM (1966) Geometric analysis of shell coiling: general problems. *J Paleontol* 40:1178–1190
- R Development Core Team (2010) R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL: <http://www.R-project.org>
- Robinson RF, Richardson CA (1998) The direct and indirect effects of suction dredging on a razor clam (*Ensis arcuatus*) population. *ICES J Mar Sci* 55:970–977
- Rohlf J (2000) TpsReg. Version 1.26. Department of Ecology and Evolution, State University of New York at Stony Brook, New York
- Rohlf J (2003a) TpsDig. Version 1.39. Department of Ecology and Evolution, State University of New York at Stony Brook, New York
- Rohlf J (2003b) TpsRel. Version 1.29. Department of Ecology and Evolution, State University of New York at Stony Brook, New York
- Rohlf FJ, Archie JW (1984) A comparison of Fourier methods for the description of wing shape in mosquitoes (Diptera: Culicidae). *Syst Zool* 33:302–317
- Roopnarine PD, Signorelli J, Laumer C (2008) Systematic, biogeographic and microhabitat -based morphometric variation of the bivalve *Anomalocardia squamosa* (Bivalvia: Veneridae: Chioninae) in Thailand. *Raffles Bull Zool* 18:95–102
- Roth VL, Mercer JM (2000) Morphometrics in development and evolution. *Am Zool* 40:801–810
- Rufino MM, Gaspar MB, Pereira AM, Vasconcelos P (2007) The use of shape to distinguish *Chamelea gallina* and *Chamelea striatula* (Bivalvia: Veneridae): linear and geometric morphometric methods. *J Morphol* 267:1433–1440
- Rufino MM, Gaspar MB, Pereira A, Maynou F, Monteiro CC (2010) Ecology of megabenthic bivalve communities from sandy beaches on the south coast of Portugal. *Sci Mar* 74:163–178
- Samadi S, Roumégoux A, Bargues MD, Mas-Coma S, Yong M, Pointier JP (2000) Morphological studies of lymnaeid snails from the human fascioliasis endemic zone of Bolivia. *J Mollus Stud* 66:31–34

- Savazzi E (1989) Burrowing mechanisms and sculptures in recent gastropods. *Lethaia* 22:31–48
- Savazzi E (1990) Biological aspects of theoretical shell morphology. *Lethaia* 23:195–212
- Savazzi E, Jefferies RPS, Signor PW (1982) Modification of the paradigm for burrowing ribs in various gastropods, crustaceans and calcichordates. *Neues Jahrb Geol Paläont Abh* 164:206–217
- Serb JM, Alejandrino A, Otárola-Castillo E, Adams DC (2011) Morphological convergence of shell shape in distantly related scallop species (Mollusca: Pectinidae). *Zool J Linn Soc* 163: 571–584
- Slice DE (1998) Morphueus et al.: software for morphometric research. Department of Ecology and Evolution, State University of New York, Stony Brook, New York
- Sousa R, Freire R, Rufino MM, Méndez J, Gaspar MB, Antunes C, Guilhermino L (2007) Genetic and shell morphological variability of the invasive bivalve *Corbicula fluminea* (Müller, 1774) in two Portuguese estuaries. *Estuar Coast Shelf Sci* 74:166–174
- Stone JR (1996) The evolution of ideas: a phylogeny of shell models. *Am Nat* 148:904–929
- Stone JR (1999) Theoretical morphology—the concept and its applications. *Science* 284:746
- Tebble N (1966) British bivalve seashells: a handbook of identification. British Museum, Edinburgh
- Tuck ID, Bailey N, Harding M, Sangster G, Howell T, Graham N, Breen M (2000) The impact of water jet dredging for razor clams, *Ensis* spp., in a shallow sandy subtidal environment. *J Sea Res* 43:65–81
- Urduy S, Goudemand N, Bucher H, Chirat R (2010a) Allometries and the morphogenesis of the molluscan shell: a quantitative and theoretical model. *J Exp Zool B Mol Dev Evol* 314:280–302
- Urduy S, Goudemand N, Bucher H, Chirat R (2010b) Growth-dependent phenotypic variation of molluscan shells: implications for allometric data interpretation. *J Exp Zool B Mol Dev Evol* 314:303–326
- Valladares A, Manríquez G, Suárez-Isla BA (2010) Shell shape variation in populations of *Mytilus chilensis* (Hupe 1854) from southern Chile: a geometric morphometric approach. *Mar Biol* 157:2731–2738
- Varela MA, González-Tizón A, Francisco-Candeira M, Martínez-Lage A (2007) Isolation and characterization of polymorphic microsatellite loci in the razor clam *Ensis siliqua*. *Mol Ecol Notes* 7:221–222